




World Health
Organization


REGIONAL OFFICE FOR Africa

Health Situation Analysis in the African Region

Atlas of Health Statistics, 2012



Health situation Analysis in the African Region Atlas of Health Statistics, 2012



AFRO Library Cataloguing-in-Publication Data

Atlas of Health Statistics of the African Region 2012

1. Health Systems
2. Health Information Systems
3. Health Policy
4. Health Services Coverage and Accessibility
5. Africa

ISBN: 978 929 023 1769 (NLM Classification: W16 and W17)

© World Health Organization. Regional Office for Africa, 2012

Publications of the World Health Organization enjoy copyright protection in accordance with the provisions of Protocol.2 of the Universal Copyright Convention. All rights reserved. Copies of this publication may be obtained from the Publication and Language Services Unit, WHO Regional Office for Africa, P.O. Box 6, Brazzaville, Republic of Congo (Tel: +47 241 39100; Fax: +47 241 39507; E-mail: afrobooks@afro.who.int). Requests for permission to reproduce or translate this publication – whether for sale or for non-commercial distribution – should be sent to the same address.

The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted lines on maps represent approximate border lines for which there may not yet be full agreement.

The mention of specific companies or of certain manufacturers' products does not imply that they are endorsed or recommended by the World Health Organization in preference to others of a similar nature that are not mentioned. Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters.

All reasonable precautions have been taken by the World Health Organization to verify the information contained in this publication. However, the published material is being distributed without warranty of any kind, either express or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall the World Health Organization or its Regional Office for Africa be liable for damages arising from its use.

Printed and bound in AFRO

More information about this publication can be obtained from:

African Health Observatory (www.aho.afro.who.int)
African Health Observatory and Knowledge Management Unit
Office of the Director, Health Systems and Services
WHO - Regional Office for Africa
Brazzaville/Republic of Congo

Contents



List of figures	iv
Message from the Regional Director	viii
Overview of the profile of the African Region	ix
1. Introduction to country context	1
2. Health status and trends	3
Life expectancy	3
Adult mortality	6
Child mortality	8
Maternal mortality	11
Age standardized death rates	12
Burden of disease	13
3. The health system	15
Health financing	15
Health workforce	23
Medical products and equipment	25
Health information	27
Health technology	28
4. Specific programmes and services	30
HIV/AIDS	30
Tuberculosis	33
Malaria	36
Immunization and vaccines development	38
Child and adolescent health	46
Maternal and newborn health	47
Gender and women's health	51
Neglected tropical diseases	54
Non-communicable diseases and conditions	55
5. Key determinants	56
Risk factors for health	56
Food safety and nutrition	57
Demography	58
Resources and infrastructure	60
Poverty and income inequality	62
Environment	63
Science and technology	66
Emergencies and disasters	67
6. Progress on MDGs	68
MDG-4: Reduce child mortality	68
MDG-5 : Improve maternal health	69
MDG-6: Combat HIV/AIDS, malaria and other diseases	71
MDG-7: Ensure environmental sustainability	74
MDG-1: Eradicate extreme poverty and hunger	75
MDG-8: Develop a global partnership for development	76
7. Explanatory notes	78

List of figures

Table : General population characteristics	ix	Figure 26 : Infant mortality rate per 1,000 live births in the African Region, 1990 and 2010	10
Figure A : Distribution of burden of diseases as % of total DALYs by broader causes (2004)	ix	Figure 27 : Annual average rate of reduction of infant mortality rate in the African Region, between 1990 and 2010	10
Figure B : Leading causes of burden of diseases (as % of total DALYs) (2004)	ix	Figure 28 : Maternal mortality ratio per 100,000 live births in the African Region, 1990 and 2010	11
Figure C : Burden of disease (% of total DALYs) by groups of disorders and conditions (2004)	ix	Figure 29 : Annual average rate of reduction of maternal mortality ratio in the African Region, between 1990 and 2010	11
Figure D : Distribution of causes of death among children aged <5 years (2010)	ix	Figure 30 : Maternal mortality ratio per 100,000 live births by WHO Region, 1990, 2000 and 2010	11
Figure E : General government expenditure on health (as % of total government expenditure)	ix	Figure 31 : Distribution of age-standardized death rates per 100,000 population in the African Region, 2008	12
Figure F : Health workforce (2000-2010)	ix	Figure 32 : Distribution of age-standardized death rates per 100,000 population due to communicable diseases in the African Region, 2008	12
Figure G : Utilization of health services (2000-2010)	ix	Figure 33 : Distribution of age-standardized death rates per 100,000 population due to non-communicable diseases in the African Region, 2008	12
Figure H : Utilization of health services (2010)	ix	Figure 34 : Distribution of age-standardized death rates per 100,000 population due to injuries and violence in the African Region, 2008	12
Figure I : Prevalence of smoking any tobacco product among adults aged > 15 years (2006)	x	Figure 35 : Distribution of burden of diseases as percentage of total DALYs by group of disorders in the African Region, 2004	13
Figure J : MDG-1: Children aged <5 years underweight	x	Figure 36 : Distribution of burden of diseases as percentage of total DALYs by broader causes, by WHO Region, 2004	13
Figure K : MDG-4: Trend in under-5 mortality rate (probability of dying by age 5; per 1,000 live births)	x	Figure 37 : Distribution of years of life lost by broader causes (%), by WHO Region, 2008	13
Figure L : MDG-5: Trend in maternal mortality ratio (per 100,000 live births)	x	Figure 38 : Leading causes of burden of diseases shown as percentage of total DALYs in the African Region, 2004	13
Figure M : MDG-5: Percentage of unmet need for family planning	x	Figure 39 : Total burden of disease in DALYs per 1,000 population by WHO Region, 2004	13
Figure N : MDG-6: Prevalence of HIV among adults aged 15-49 years, by Region (2009)	x	Figure 40 : Distribution of burden of diseases as percentage of total DALYs by broader causes in the African Region, 2004	14
Figure O : MDG-6: Percentage of population living with advanced HIV infection who are receiving antiretroviral therapy, by Region (2010)	x	Figure 41 : Distribution of years of life lost by broader causes (%) in the African Region, 2008	14
Figure P : MDG-6: Malaria mortality rate per 100,000 population, by Region (2008)	x	Figure 42 : Total expenditure on health as percentage of GDP in the African Region, 2010	15
Figure Q : MDG-6: Prevalence of tuberculosis (per 100,000 population), by Region (2009)	x	Figure 43 : Growth rate of total expenditure on health as percentage of GDP in the African Region, 2000 and 2010	15
Figure R : MDG-7: Population using improved drinking water sources	x	Figure 44 : Trend in total expenditure on health as percentage of GDP in the African Region, 2004 to 2009	15
Figure S : MDG-7: Population using improved sanitation	x	Figure 45 : Total expenditure on health as percentage of GDP by WHO Region, 2000 and 2009	15
Figure 1 : WHO Regions	1	Figure 46 : Per capita total expenditure on health (PPP int. \$) in the African Region, 2000 and 2010	16
Figure 2 : Population size (in thousands) of countries of the African Region, 2010	1	Figure 47 : Per capita total expenditure on health (PPP int. \$) by WHO Region, 2000 and 2009	16
Figure 3 : Age distribution of population by WHO Region, 2010	1	Figure 48 : Per capita total expenditure on health (PPP int. \$) in the African Region, 2010	16
Figure 4 : Population size (in percentage) by WHO Region, 2010	2	Figure 49 : Trend in average per capita total expenditure on health (PPP int. \$) in the African Region, 2004 to 2009	16
Figure 5 : Age distribution of the population in the African Region, 2010	2	Figure 50 : General government expenditure on health as percentage of total government expenditure in the African Region, 2000 and 2010	17
Figure 6 : Annual growth rate (%) distribution of population by WHO Region, 2000-2010	2	Figure 51 : General government expenditure on health as percentage of total government expenditure by WHO Region, 2000 and 2009	17
Figure 7 : Life expectancy at birth in years in the African Region, 2009	3	Figure 52 : General government expenditure on health as percentage of total government expenditure in the African Region, 2010	17
Figure 8 : Life expectancy at birth in years by WHO Region, 1990 and 2009	3	Figure 53 : Trend in general government expenditure on health as percentage of total government expenditure in the African Region, 2004 to 2009	17
Figure 9 : Life expectancy at birth in years by WHO Region, by sex, 2009	3	Figure 54 : General government expenditure on health as percentage of total expenditure on health in the African Region, 2000 and 2010	18
Figure 10 : Life expectancy at birth in years in the African Region, 1990 and 2009	4	Figure 55 : General government expenditure on health as percentage of total expenditure on health by WHO Region, 2000 and 2009	18
Figure 11 : Life expectancy at birth in years in the African Region, by sex, 2009	4	Figure 56 : General government expenditure on health as percentage of total expenditure on health in the African Region, 2010	18
Figure 12 : Healthy life expectancy at birth in years by WHO Region, by sex, 2007	5	Figure 57 : Trend in general government expenditure on health as percentage of total expenditure on health in the African Region, 2004 to 2009	18
Figure 13 : Healthy life expectancy at birth in years in the African Region, by sex, 2007	5	Figure 58 : ExTERNAL resources for health as percentage of total expenditure on health in the African Region, 2000 and 2010	19
Figure 14 : Trend in life expectancy at birth in years in the African Region, by sex, 2003 to 2009	5	Figure 59 : ExTERNAL resources for health as percentage of total expenditure on health by WHO Region, 2000 and 2009	19
Figure 15 : Adult mortality rate per 1,000 population in the African Region, 2009	6	Figure 60 : ExTERNAL resources for health as percentage of total expenditure on health in the African Region, 2010	19
Figure 16 : Adult mortality rate per 1,000 population in the African Region, by sex, 2009	6		
Figure 17 : Adult mortality rate per 1,000 population by WHO Region, 1990 and 2009	6		
Figure 18 : Adult mortality rate per 1,000 population by WHO Region, by sex, 2009	6		
Figure 19 : Adult mortality rate per 1,000 population in the African Region, 1990 and 2009	7		
Figure 20 : Annual average rate of reduction of Adult mortality rate in the African Region, between 1990 and 2009	7		
Figure 21 : Under-5 mortality rate per 1,000 live births in the African Region, 2010	8		
Figure 22 : Under-5 mortality rate per 1,000 live births by WHO Region, both sexes, 1990, 2000 and 2010	8		
Figure 23 : Infant mortality rate per 1,000 live births by WHO Region, 1990, 2000 and 2010	8		
Figure 24 : Under-5 mortality rate per 1,000 live births in the African Region, 1990 and 2010	9		
Figure 25 : Annual average rate of reduction of under-five mortality in the African Region, between 1990 and 2010	9		

Figure 61: Trend in external resources for health as percentage of total expenditure on health in the African Region, 2004 to 2009	19	Figure 97: Availability of national standards or recommended lists of medical devices in the African Region, 2010	29
Figure 62: Out-of-pocket expenditure as percentage of private expenditure on health in the African Region, 2000 and 2010	20	Figure 98: Presence of types of lists recommending health technology in the African Region, 2010	29
Figure 63: Out-of-pocket expenditure as percentage of private expenditure on health by WHO Region, 2000 and 2009	20	Figure 99: HIV/AIDS mortality rate (per 100 000 population) in the African Region, 2009	30
Figure 64: Out-of-pocket expenditure as percentage of private expenditure on health in the African Region, 2010	20	Figure 100: HIV/AIDS mortality rate (per 100 000 population) by WHO Region, 2000 and 2009	30
Figure 65: Trend in out-of-pocket expenditure as percentage of private expenditure on health in the African Region, 2004 to 2009	20	Figure 101: Prevalence of HIV (per 100 000 population) by WHO Region, 2000 and 2009	30
Figure 66: Private expenditure on health as percentage of total expenditure on health in the African Region, 2000 and 2010	21	Figure 102: Prevalence of HIV (per 100 000 population) in the African Region, 2000 and 2009	30
Figure 67: Private expenditure on health as percentage of total expenditure on health by WHO Region, 2000 and 2009	21	Figure 103: Percentage of people with advanced HIV infection receiving antiretroviral (ARV) combination therapy in the African Region, 2010	31
Figure 68: Private expenditure on health as percentage of total expenditure on health in the African Region, 2010	21	Figure 104: Percentage of antiretroviral therapy among people with HIV/AIDS in need of treatment by WHO Region, 2007 and 2010	31
Figure 69: Trend in private expenditure on health as percentage of total expenditure on health in the African Region, 2004 to 2009	21	Figure 105: Trend in antiretroviral therapy coverage (%) among HIV-infected pregnant women for PMTCT in the African Region, 2005 to 2010	31
Figure 70: Per capita government expenditure on health (PPP int. \$) in the African Region, 2000 and 2010	22	Figure 106: Percentage of people receiving antiretroviral therapy in the African Region, 2007 and 2010	31
Figure 71: Per capita government expenditure on health (PPP int. \$) by WHO Region, 2000 and 2009	22	Figure 107: Prevalence of condom use by adults aged 15-49 years during higher-risk sex in the African Region, 2005-2009	32
Figure 72: Per capita government expenditure on health (PPP int. \$) in the African Region, 2010	22	Figure 108: Population aged 15-24 years of age with comprehensive knowledge of HIV/AIDS in the African Region, 2005-2009	32
Figure 73: Trend per capita government expenditure on health (PPP int. \$) in the African Region, 2004 to 2009	22	Figure 109: Tuberculosis death rate per 100,000 population per year in the African Region, by country, 2000 and 2007	33
Figure 74: Physician-to-population ratio (per 10,000 population) in the African Region, 2005-2010	23	Figure 110: Tuberculosis incidence rate per 100,000 population per year in the African Region, by country, 2000 and 2010	33
Figure 75: Physician-to-population ratio (per 10,000 population) by WHO Region, 2005-2010	23	Figure 111: Tuberculosis prevalence per 100,000 population per year in the African Region, by country, 2010	34
Figure 76: Nursing and midwifery personnel-to-population ratio (per 10,000 population) in the African Region, 2005-2010	23	Figure 112: Tuberculosis prevalence per 100,000 population per year by WHO Region, 2000 and 2010	34
Figure 77: Nursing and midwifery personnel-to-population ratio (per 10,000 population) by WHO Region, 2005-2010	23	Figure 113: Trend in tuberculosis prevalence per 100,000 population per year in the African Region, by country, 2004 and 2010	34
Figure 78: Dentistry personnel density (per 10,000 population) in the African Region, 2000-2010	24	Figure 114: Tuberculosis prevalence per 100,000 population per year in the African Region, 2000 and 2010	34
Figure 79: Dentistry personnel density (per 10,000 population) by WHO Region, 2005-2010	24	Figure 115: Percentage of tuberculosis detection rate under in the African Region, 2007	35
Figure 80: Pharmaceutical personnel density (per 10,000 population) in the African Region, 2005-2010	24	Figure 116: Percentage of tuberculosis detection rate under DOTS by WHO Region, 2000 and 2007	35
Figure 81: Pharmaceutical personnel density (per 10,000 population) by WHO Region, 2005-2010	24	Figure 117: Trend in percentage of tuberculosis detection rate under DOTS in the African Region, by country, 2003 to 2007	35
Figure 82: Median percentage availability of selected generic medicines in a sample of health facilities in the African Region, countries with data in 2001-2009 (Private, Public)	25	Figure 118: Percentage of tuberculosis detection rate under DOTS in the African Region, by country, 2000 and 2007	35
Figure 83: Median consumer price ratio of selected medicines (ratio of median local unit price to management sciences for health international reference price) in the African Region, countries with data in 2001-2009 (Private, Public)	25	Figure 119: Percentage of children under 5 years of age sleeping under insecticide-treated bed nets in the African Region, by country, 2005-2009	36
Figure 84: Hospital beds per 10,000 population in the African Region, 2005-2011	26	Figure 120: Proportion of children under 5 years of age with fever being treated with antimalarial drugs in the African Region, by country, 2005-2010	36
Figure 85: Hospital beds per 10,000 population by WHO Region, 2005-2011	26	Figure 121: Malaria mortality rate in the African Region, 2008	37
Figure 86: Psychiatric beds (per 10,000 population) in the African Region, 2005-2011	26	Figure 122: Malaria mortality rate by WHO Region, 2006 and 2008	37
Figure 87: Psychiatric beds (per 10,000 population) in the WHO Region, 2005-2011	26	Figure 123: Trend in reported clinical malaria cases (in millions) in the African Region, 1990 to 2009	37
Figure 88: Radiotherapy units per 1,000,000 population in the African Region, 2010	27	Figure 124: Reported cases of malaria (in thousands) in the African Region, 2010	37
Figure 89: Percentage of civil registration coverage for deaths in the African Region, 2000-2009	27	Figure 125: Percentage of neonates protected at birth against neonatal tetanus in the African Region, 2011	38
Figure 90: Percentage of civil registration coverage for births in the African Region, 2005-2010	27	Figure 126: Percentage of neonates protected at birth against neonatal tetanus by WHO Region, 1990 and 2011	38
Figure 91: Availability of national list of approved medical devices for procurement or reimbursement, African Region, 2010	28	Figure 127: Trend in percentage of neonates protected at birth against neonatal tetanus in the African Region, 1980 to 2011	38
Figure 92: Availability of technical specifications of medical devices to support procurement donations in the African Region, 2010	28	Figure 128: Percentage of neonates protected at birth against neonatal tetanus in the African Region, 1990 and 2011	38
Figure 93: Presence of health technology (medical device) national policy, Africa Region, 2010	28	Figure 129: Percentage immunization coverage among 1-year-olds for DTP3 in the African Region, 2011	39
Figure 94: Presence of national guidelines, policies or recommendations on the procurement of medical devices, African Region, 2010	28	Figure 130: Percentage immunization coverage among 1-year-olds for DTP3 by WHO Region, 1990 and 2011	39
Figure 95: Procurement of medical devices carried out at the national level, African Region, 2010	29	Figure 131: Trend in percentage immunization coverage among 1-year-olds for DTP3 in the African Region, 1980 to 2011	39
Figure 96: Presence of units in the ministries of health responsible for the implementation of the health technology national policy in the African Region, 2010	29	Figure 132: Percentage immunization coverage among 1-year-olds for DTP3 in the African Region, by country, 1990 and 2011	39
		Figure 133: Percentage immunization coverage among 1-year-olds for HepB3 in the African Region, 2011	40

Figure 134 : Percentage immunization coverage among 1-year-olds for HepB3 by WHO Region, 1990 and 2011	40	Figure 172 : Distribution of main causes of antenatal death in the African Region, 2004	50
Figure 135 : Trend in immunization coverage among 1-year-olds for HepB3 in the African Region, 1997 to 2011	40	Figure 173 : Percentage of births by caesarean section in the African Region, 2005-2010	50
Figure 136 : Percentage immunization coverage among 1-year-olds for HepB3 in the African Region, 2000 and 2011	40	Figure 174 : Adolescent fertility rate (per 1000 girls, 15-19 years of age) by WHO Region, 2005-2010	51
Figure 137 : Percentage immunization coverage among 1-year-olds for BCG in the African Region, 2011	41	Figure 175 : Total women of childbearing age (15-49 years), in thousands, in the African Region, 2010	51
Figure 138 : Percentage immunization coverage among 1-year-olds for BCG by WHO Region, 1990 and 2011	41	Figure 176 : Adolescent fertility rate (per 1000 girls, 15-19 years of age) in the African Region, 2005-2010	51
Figure 139 : Trend in percentage immunization coverage among 1-year-olds for BCG in the African Region, 1980 to 2011	41	Figure 177 : Contraceptive prevalence in the African Region, 2005-2010	52
Figure 140 : Percentage immunization coverage among 1-year-olds for BCG in the African Region, 1990 and 2011	41	Figure 178 : Contraceptive prevalence by WHO Region, 2005-2010	52
Figure 141 : Percentage immunization coverage among 1-year-olds for Hib in the African Region, 2010	42	Figure 179 : Percent of contraceptive use (modern method) among married women 15-49 years old in the African Region, 2000-2009	52
Figure 142 : Percentage immunization coverage among 1-year-olds for Hib by WHO Region, 1990 and 2011	42	Figure 180 : Contraceptive prevalence in the African Region, 2005-2010	52
Figure 143 : Trend in percentage immunization coverage among 1-year-olds for Hib in the African Region, 2000 to 2011	42	Figure 181 : Total fertility rate per woman in the African Region, 2010	53
Figure 144 : Percentage immunization coverage among 1-year-olds for Hib in the African Region, 1990 and 2011	42	Figure 182 : Total fertility rate per woman by WHO Region, 1990 and 2010	53
Figure 145 : Percentage immunization coverage among 1-year-olds for Measles in the African Region, 2010	43	Figure 183 : Trend in total fertility rate per woman in the African Region, 2004 to 2010	53
Figure 146 : Percentage immunization coverage among 1-year-olds for Measles by WHO Region, 1990 and 2011	43	Figure 184 : Total fertility rate per woman in the African Region, 1990 and 2010	53
Figure 147 : Trend in percentage immunization coverage among 1-year-olds for Measles in the African Region, 1980 to 2011	43	Figure 185 : Number of reported cases of leprosy in the African Region, by country, 2008	54
Figure 148 : Percentage immunization coverage among 1-year-olds for Measles in the African Region, 1990 and 2011	43	Figure 186 : Number of reported cases of leprosy by WHO Region, 2010	54
Figure 149 : Percentage immunization coverage among 1-year-olds for PCV in the African Region, 2010	44	Figure 187 : Trend in number of reported cases of leprosy (in thousands) in the African Region, 2005 to 2010	54
Figure 150 : Percentage immunization coverage among 1-year-olds for PCV by WHO Region, 1990 and 2011	44	Figure 188 : Number of reported cases of leprosy in the African Region, 2007 and 2010	54
Figure 151 : Trend in percentage immunization coverage among 1-year-olds for PCV in the African Region, 1980 to 2011	44	Figure 189 : Distribution of causes of non-communicable burden of diseases (percentage of total DALYs) in the African Region, 2004	55
Figure 152 : Percentage immunization coverage among 1-year-olds for PCV in the African Region, 1990 and 2011	44	Figure 190 : Distribution of causes of neuropsychiatric burden of diseases (percentage of total DALYs) in the African Region, 2004	55
Figure 153 : Percentage immunization coverage among 1-year-olds for Polio in the African Region, 2010	45	Figure 191 : Distribution of causes of intentional and non-intentional injuries (percentage of total DALYs) in the African Region, 2004	55
Figure 154 : Percentage immunization coverage among 1-year-olds for Hib by WHO Region, 1990 and 2011	45	Figure 192 : Insufficiently active in persons aged 15 years or older, by sex, in the African Region, 2008	55
Figure 155 : Trend in percentage immunization coverage among 1-year-olds for Polio in the African Region, 2000 to 2011	45	Figure 193 : Percentage of current tobacco use in persons 15 years of age or older by WHO Region, 2009	56
Figure 156 : Percentage immunization coverage among 1-year-olds for Polio in the African Region, 1990 and 2011	45	Figure 194 : Percentage of current tobacco use in persons 15 years of age or older in the African Region, 2009	56
Figure 157 : Distribution of causes of death among children aged under 5 years (in percentage) in the African Region, 2010	46	Figure 195 : Alcohol consumption (litres per person) among adults aged 15 years of age or older by WHO Region, 2005	56
Figure 158 : Percentage of infants exclusively breastfed for the first 6 months of life in the African Region, 2005-2011	46	Figure 196 : Alcohol consumption (litres per person) among adults aged 15 years of age or older in the African Region, 2005	56
Figure 159 : Percentage of infants exclusively breastfed for the first 6 months of life by WHO Region, 2005-2011	46	Figure 197 : Percentage of underweight children under 5 years of age in the African Region, 1990-1995 and 2005-20011	57
Figure 160 : Percentage of infants exclusively breastfed for the first 6 months of life in the African Region, 2005-2011	46	Figure 198 : Percentage of underweight children under 5 years of age in the African Region, 2005-20011	57
Figure 161 : Percentage of low-birthweight newborns by WHO Region, 2005-2010	47	Figure 199 : Percentage of stunted children under 5 years of age in the African Region, 2005-20011	57
Figure 162 : Percentage of low-birthweight newborns in the African Region, 2005-2010	47	Figure 200 : Percentage of overweight children under 5 years of age in the African Region, 2005-20011	57
Figure 163 : Malnutrition prevalence, weight for age (% of children under 5) in the African Region, 2000-2009	47	Figure 201 : Annual growth rate (in percent) of population in the African Region, 1989-1999 and 2000-2010	58
Figure 164 : Percentage of births attended by skilled health personnel in the African Region, 2005-2011	48	Figure 202 : Annual growth rate (in percent) of population by WHO Region, 1989-1999 and 2000-2010	58
Figure 165 : Percentage of births attended by skilled health personnel by WHO Region, 1990-1999 and 2005-2011	48	Figure 203 : Percentage of population living in urban areas in the African Region, 2010	58
Figure 166 : Estimated workforce shortage to attain 95% skilled birth attendance by 2015 in the African Region, 2010	48	Figure 204 : Trend in percentage of population living in urban areas in the African Region, 2005 to 2010	58
Figure 167 : Percentage of births attended by skilled health personnel in the African Region, by country, 1990-1999 and 2005-2011	48	Figure 205 : Total fertility rate per woman in the African Region, 1990 and 2010	59
Figure 168 : Percentage of antenatal care coverage, at least one visit, in the African Region, 2005-2011	49	Figure 206 : Total fertility rate per woman, by WHO Region, 1990 and 2010	59
Figure 169 : Percentage of antenatal care coverage, at least four visits, in the African Region, 2005-2011	49	Figure 207 : Total fertility rate per woman in the African Region, 2010	59
Figure 170 : Percentage of births by caesarean section in the African Region, 2005-2010	50	Figure 208 : Trend in total fertility rate per woman in the African Region, 2000 to 2010	59
Figure 171 : Percentage of births by caesarean section by WHO Region, 2005-2010	50	Figure 209 : Gross national income per capita (PPP int. \$) in the African Region, 1990 and 2010	60
		Figure 210 : Gross national income per capita (PPP int. \$) by WHO Region, 1990 and 2010	60
		Figure 211 : Gross national income per capita (PPP int. \$) in the African Region, 2010	60
		Figure 212 : Trend in gross national income per capita (PPP int. \$) in the African Region, 2003 to 2010	60
		Figure 213 : Electrification rate in the African Region, 2009	61
		Figure 214 : Paved roads as percentage of all roads in the African Region, 2000-2007	61

Figure 215: Electrification rate in the African Region, 2000-2005	61	Figure 240: Percentage of births attended by skilled health personnel in the African Region, by country, 2005-2011 and the MDG Target 2015	70
Figure 216: Percentage of the population living under \$1 (PPP int. \$) a day (i.e. absolute poverty) in the African Region, 2000-2009	62	Figure 241: Percentage of unmet need for family planning in the African Region, by country, 1990-1999 and 2005-2010	70
Figure 217: Percentage of the population living under \$1 (PPP int. \$) a day (i.e. absolute poverty) by WHO Region, 2000-2009	62	Figure 242: Percentage of children under 5 years of age sleeping under insecticide-treated bed nets in the African Region, 2000-2004 and 2005-2009	71
Figure 218: Share of income by poorest and richest section of the population in the African Region, 1989-2005	62	Figure 243: Percentage of children under 5 years of age with fever being treated with antimalarial drugs in the African Region, 2000-2004 and 2005-2010	71
Figure 219: Percentage of population using an improved drinking water source in the African Region, 1990 and 2010	63	Figure 244: Malaria mortality rate (per 100,000 population) in the African Region, by country, 2008	71
Figure 220: Percentage of urban and rural population with access to improved drinking-water source in the African Region, 2010	63	Figure 245: Trend in death rate associated with malaria in African Region, 1990-2009	71
Figure 221: Percentage of population using an improved drinking water source by WHO Region, 1990 and 2008	64	Figure 246: Percentage of pregnant women living with HIV infection who are receiving antiretroviral therapy for preventing mother-to-child transmission, 2010	72
Figure 222: Percentage of urban and rural population with access to improved drinking-water source by WHO Region, 2008	64	Figure 247: Percentage of people 15-49 years of age living with HIV in the African Region, by country, 2009	72
Figure 223: Percentage of population using an improved sanitation facilities by WHO Region, 1990 and 2008	64	Figure 248: Trend in contraceptive prevalence rate in African Region, 2000-2010	72
Figure 224: Percentage of the population using improved sanitation facilities by WHO Region, urban and rural, 2008	64	Figure 249: Prevalence of tuberculosis (per 100,000 population) in the African Region, by country, 2009	73
Figure 225: Trend in percentage of the urban and rural population with access to improved drinking-water source in the African Region, 2000 to 2008	64	Figure 250: Percentage of population living with advanced HIV infection who are receiving antiretroviral therapy, 2010	73
Figure 226: Trend in percentage of population using improved sanitation facilities in the African Region, urban and rural, 2000 to 2008	64	Figure 251: Trend in Tuberculosis case detection rate (% , all forms) in Sub-Saharan Africa, 1990 - 2010	73
Figure 227: Percentage of population using improved sanitation facilities in the African Region, 1990 and 2010	65	Figure 252: Percentage of the population using improved drinking water sources in the African Region, by country, 2010 and the MDG target 2015	74
Figure 228: Percentage of the population using improved sanitation facilities in the African Region, urban and rural, 2010	65	Figure 253: Percentage of the population using improved sanitation facilities in the African Region, by country, 2010 and the MDG target 2015	74
Figure 229: Population who are telephone fixed subscribers (per 10,000) in the African Region, 2009	66	Figure 254: Percentage of underweight children under 5 years of age in the African Region, by country, 2005-2011 and the MDG target	75
Figure 230: Population who are internet users (per 10,000) in the African Region, 2008	66	Figure 255: Percentage of children aged under 5 years that are underweight in the African Region, by country, 2005-2011	75
Figure 231: Under-five mortality by WHO Region, both sexes, 1990 and 2009	67	Figure 256: Annual average rate of reduction of underweight children under 5 years of age in the African Region, by country, 1990-1995 and 2005-2011	75
Figure 232: The total number of internally displaced people in the African Region, 2001 and 2009	67	Figure 257: Official development assistance (ODA) received as percentage of GDP in the African Region, by country, 2010	76
Figure 233: The total number of refugees (in thousands) in the African Region, by country of origin, 2010	67	Figure 258: Total debt service as percentage of exports of goods, services and income in the African Region, by country, 1990 and 2009	76
Figure 234: Under-5 mortality rate (per 1000 live births) in WHO Regions, both sexes, 1990 and 2010	68	Figure 259: Trade (% of GDP) in Sub-Saharan Africa, 1990-2010	76
Figure 235: Under-5 mortality rate (per 1000 live births) in the African Region, by country, 2010 and the MDG target	68	Figure 260: Percentage of the population who are internet users in the African Region, by country, 2010	77
Figure 236: Annual average rate of reduction of under-5 mortality in the African Region, 1990 and 2010	68	Figure 261: Cellular or mobiles (per 100 population) in the African Region, by country, 2010	77
Figure 237: Maternal mortality ratio (per 100 000 live births) in WHO Regions in 2010	69	Figure 262: Unemployment, youth total (% of total labor force ages 15-24) in the African Region, by country, 1990-1999 and 2000-2009	77
Figure 238: Maternal mortality ratio (per 100 000 live births) in the African Region, by country, 2010 and the MDG target	69		
Figure 239: Annual average rate of reduction of maternal mortality (per 100 000 live births) in the African Region, by country, 1990 and 2010	69		

Message from the Regional Director



The *Atlas of Health Statistics, 2012*, which provides a health situation analysis of WHO's African Region, is the most significant data output of the African Health Observatory (www.aho.afro.who.int). Now in its second edition, the *Atlas* is building on the ground-breaking work that was carried out in preparing the initial edition. Not only has it been updated for 2012, but its coverage has expanded and further indicators have been included. Another new development is the presence of the *Atlas* on the African Health Observatory web portal. It is being launched not merely as an electronic document, but as interactive web pages within the Observatory, allowing users to carry out searches and conduct analyses of their own. We aim to develop the *Atlas* on an ongoing basis, expanding its reach, indicators and accuracy as we go.

Of course all the data comes from the countries, and we are entirely reliant on data collection, cleaning, correction, evaluation and assessment carried out first of all at country level in each of the 46 Member Countries of WHO's African Region. These data are further reviewed and refined in WHO, both in its African country offices and Regional Office, and by technical experts at WHO headquarters in Geneva. Mortality estimates that are used to monitor internationally agreed goals, such as the MDGs, are produced by inter-agency groups consisting of members from WHO, UNICEF, and World Bank among others. The results of this system of analysis is data which is as good as can be extracted from the raw inputs.

Looking back to the raw inputs, however, it is clear that the quality, quantity, frequency of collection, and timeliness of data depends very much on the strength of the national health information systems, which include data collection at the district and peripheral levels. With some notable exceptions, this has been an area of weakness within most national health systems. By and large, the development of national health information systems has been slow and uneven, despite many efforts over the years.

WHO seeks to support countries in strengthening their national health information systems, and one mechanism that is being developed in response to demands from the countries is the establishment of a network of national health observatories. With support from WHO's Regional Office for Africa, a number of countries have taken steps to set up such observatories, often with direct links to the district level, as a way to reinforce the national health information system. The national observatories also link to the African Health Observatory, in a collaborative, two-way system of information, evidence and knowledge exchange. Such observatories serve at both the regional and national levels as platforms for other activities designed to foster monitoring and evaluation, which are essential components of the cycle of development and policy work that lead to national health policies and health development plans.

Thus, the collaborative networking approach embraced by the African Health Observatory and the national observatories is intended to provide a continuum between work at the regional and national levels, offering a platform for many disparate supporting mechanisms and methodologies. This should lead to a marked decrease of the fragmentation of efforts so frequently found in public health policy and development work. The *Atlas* is a product and promoter of such collaborative networking.

Overview of the Profile of the African Region



All data sources are given in the African Health Observatory at www.aho.afro.who.int

... No data

The WHO African Region



Table : General population characteristics

		Africa	Global
Population size (in thousands)	2009	824,401	6,816,573
	2010	836,970	6,860,318
Population - Living in urban areas (%)	1990	29	43
	2009	38	50
Life expectancy at birth (years)	1990	51	64
	2009	54	68
Adult mortality rate (probability of dying between 15 and 60 years per 1000 population)	1990	366	207
	2009	383	176
Per capita total expenditure on health (PPP int. \$)	2000	87	566
	2009	157	990
Gross national income per capita (PPP int. \$)	1990	1,319	4,862
	2008	2,279	10,290

Figure A : Distribution of burden of diseases as % of total DALYs by broader causes (2004)




Figure B : Leading causes of burden of diseases (as % of total DALYs) (2004)




Figure C : Burden of disease (% of total DALYs) by groups of disorders and conditions (2004)




Figure D : Distribution of causes of death among children aged <5 years (2010)




Figure E : General government expenditure on health (as % of total government expenditure)




Figure F : Health workforce, 2005–2010




Figure G : Utilization of health services




Figure H : Utilization of health services, 2010





Figure I : Prevalence of smoking any tobacco product among adults aged > 15 years, 2009



Overview of Progress on the MDGs in the African Region



Figure J : MDG-1: Children aged < 5 years underweight




Figure K : MDG-4: Trend in under-5 mortality rate (probability of dying by age 5; per 1,000 live births)




Figure L : MDG-5: Trend in maternal mortality ratio (per 100,000 live births)




Figure M : MDG-5: Percentage of unmet need for family planning




Figure N : MDG-6: Prevalence of HIV among adults aged 15–49 years, by Region, 2009




Figure O : MDG-6: Percentage of population living with advanced HIV infection who are receiving antiretroviral therapy, by Region, 2010




Figure P : MDG-6: Malaria mortality rate per 100,000 population, by Region, 2008




Figure Q : MDG-6: Prevalence of tuberculosis (per 100,000 population), by Region, 2010




Figure R : MDG-7: Population using improved drinking water sources





Figure S : MDG-7: Population using improved sanitation



1. Introduction to country context

Figure 1 : WHO Regions

- African Region
- Eastern Mediterranean Region
- European Region
- Region of the Americas
- South-East Asia Region
- Western Pacific Region



Figure 2 : Population size (in thousands) of countries of the African Region, 2010




Figure 3 : Age distribution of population by WHO Region, 2010




Figure 4 : Population size (in percentage) by WHO Region, 2010





Figure 6: Annual growth rate (%) distribution of population by WHO Region, 2000-2010



■ Aged over 60 (%)
■ Aged 15 - 59 (%)
■ Aged under 15 (%)


Figure 5 : Age distribution of the population in the African Region, 2010

Country	Aged under 15 (%)	Aged 15 - 59 (%)	Aged over 60 (%)
Niger	49	47	4
Uganda	48	48	4
Mali	47	49	4
Angola	47	49	4
Malawi	46	49	5
Zambia	46	49	5
DR Congo	46	50	4
Tanzania	45	50	5
Burkina Faso	45	51	4
Chad	45	50	5
Gambia	44	53	3
Benin	44	51	5
Mozambique	44	51	5
Senegal	44	52	4
Comoros	43	53	4
Rwanda	43	53	4
Guinea	43	52	5
Liberia	43	53	4
Madagascar	43	52	5
Nigeria	43	52	5
Sierra Leone	43	53	4
Eritrea	42	54	4
Kenya	42	54	4
Congo	41	53	6
Cameroon	41	54	5
Côte d'Ivoire	41	53	6
Guinea-Bissau	41	54	5
Ethiopia	41	54	5
Mauritania	40	56	4
Togo	40	55	5
Central African Republic	40	54	6
Sao Tome and Principe	40	55	5
Ghana	39	55	6
Zimbabwe	39	55	6
Equatorial Guinea	39	57	4
Burundi	38	57	5
Swaziland	38	57	5
Lesotho	37	57	6
Namibia	36	58	6
Gabon	35	59	6
Botswana	33	61	6
Cape Verde	32	61	7
South Africa	30	63	7
Algeria	27	66	7
Seychelles	23	66	11
Mauritius	22	67	11

2. Health Status and Trends


Life expectancy

Figure 7 : Life expectancy at birth in years in the African Region, 2009




1990
2009

Figure 8 : Life expectancy at birth in years by WHO Region, 1990 and 2009



Female
Male

Figure 9 : Life expectancy at birth in years by WHO Region, by sex, 2009





Life expectancy

1990
2009

Figure 10: Life expectancy at birth in years in the African Region, 1990 and 2009





Figure 11: Life expectancy at birth in years in the African Region, by sex, 2009

Female
Male



Life expectancy

Figure 12 : Healthy life expectancy* at birth in years by WHO Region, by sex, 2007




Figure 13 : Healthy life expectancy at birth in years in the African Region, by sex, 2007





Figure 14: Trend in life expectancy at birth in years in the African Region, by sex, 2003 to 2009





Adult mortality

Figure 15 : Adult mortality rate per 1,000 population in the African Region, 2009





Figure 17 : Adult mortality rate per 1,000 population by WHO Region, 1990 and 2009




Figure 18 : Adult mortality rate per 1,000 population by WHO Region, by sex, 2009





Figure 16 : Adult mortality rate per 1,000 population in the African Region, by sex, 2009



Adult mortality

Figure 19 : Adult mortality rate per 1,000 population in the African Region, 1990 and 2009





Figure 20 : Annual average rate of reduction of adult mortality rate in the African Region, between 1990 and 2009



Child mortality

Figure 21: Under-5 mortality rate per 1,000 live births in the African Region, 2010




Figure 22 : Under-5 mortality rate per 1,000 live births by WHO Region, 1990, 2000 and 2010





Figure 23 : Infant mortality rate per 1,000 live births by WHO Region, 1990, 2000 and 2010



Child mortality

1990
2010

Figure 24 : Under-5 mortality rate per 1,000 live births in the African Region, 1990 and 2010





Figure 25 : Annual average rate of reduction of under-five mortality in the African Region, between 1990 and 2010



Child mortality

1990
2010

Figure 26 : Infant mortality rate per 1,000 live births in the African Region, 1990 and 2010






Figure 27 : Annual average rate of reduction of infant mortality rate in the African Region, between 1990 and 2010



Maternal mortality


1990
2010

Figure 28: Maternal mortality ratio per 100,000 live births in the African Region, 1990 and 2010



Countries of the African Region without data are not included in the chart.


Figure 29: Annual average rate of reduction of maternal mortality ratio in the African Region, between 1990 and 2010



Countries of the African Region without data are not included in the chart.

Africa
South-East Asia
Eastern Mediterranean
Global
Western Pacific
Americas
Europe

Figure 30: Maternal mortality ratio per 100,000 live births by WHO Region, 1990 to 2010



Age standardized death rates

■ Injuries
■ Noncommunicable
■ Communicable

Figure 31 : Distribution of age-standardized death rates* per 100,000 population in the African Region, 2008

Country	Communicable	Noncommunicable	Injuries
Africa	798	779	107
Zimbabwe	1,552	622	73
Lesotho	1,255	774	141
Swaziland	1,200	867	208
Malawi	1,156	999	189
Central African Republic	1,060	870	151
Sierra Leone	1,042	763	92
Chad	1,009	866	119
South Africa	983	635	72
Zambia	961	938	176
Mozambique	957	908	153
Burundi	943	839	124
DR Congo	932	837	185
Guinea-Bissau	928	916	110
Côte d'Ivoire	904	942	172
Cameroon	861	879	111
Nigeria	832	809	76
Mali	827	733	72
Angola	819	842	112
Uganda	810	888	178
Burkina Faso	801	810	108
Tanzania	782	745	120
Liberia	782	766	63
Guinea	759	932	121
Botswana	739	606	107
Niger	730	647	44
Ethiopia	721	903	139
Equatorial Guinea	720	854	117
Congo	692	811	140
Namibia	670	791	160
Togo	635	716	63
Kenya	624	681	116
Benin	618	804	87
Ghana	608	711	91
Rwanda	595	740	92
Mauritania	575	746	88
Gabon	545	660	82
Gambia	525	735	72
Senegal	509	665	58
Comoros	472	789	76
Madagascar	408	706	58
Eritrea	303	670	92
Sao Tome and Principe	279	605	45
Cape Verde	213	538	71
Algeria	202	523	47
Seychelles	149	591	48
Mauritius	62	664	42

Figure 32 : Distribution of age-standardized death rates* per 100,000 population due to communicable diseases in the African Region, 2008




Figure 33 : Distribution of age-standardized death rates* per 100,000 population due to non-communicable diseases in the African Region, 2008





Figure 34 : Distribution of age-standardized death rates* per 100,000 population due to injuries and violence in the African Region, 2008



Burden of disease

Figure 35 : Distribution of burden of diseases as percentage of total DALYs by group of disorders in the African Region, 2004




Figure 38 : Leading causes of burden of diseases shown as percentage of total DALYs in the African Region, 2004




Figure 36 : Distribution of burden of diseases as percentage of total DALYs by broader causes, by WHO Region, 2004




Figure 37 : Distribution of years of life lost* by broader causes (%), by WHO Region, 2008






Figure 39 : Total burden of disease in DALYs* per 1,000 population by WHO Region, 2004



*The disability-adjusted life-year (DALY) provides a consistent and comparative description of the burden of diseases and injuries needed to assess the comparative importance of diseases and injuries in causing premature death, loss of health and disability in different populations. The DALY extends the concept of potential years of life lost due to premature death to include equivalent years of 'healthy' life lost by virtue of being in states of poor health or disability. One DALY can be thought of as one lost year of 'healthy' life, and the burden of disease can be thought of as a measurement of the gap between current health status and an ideal situation where everyone lives into old age, free of disease and disability. WHO. Burden of Diseases Update 2004. Geneva, July 2008.


Burden of disease

Figure 40 : Distribution of burden of diseases as percentage of total DALYs by broader causes in the African Region, 2004



■ Injuries
■ Noncommunicable
■ Communicable

Figure 41 : Distribution of years of life lost by broader causes (%) in the African Region, 2008




■ Injuries
■ Noncommunicable
■ Communicable

3. The health system


Health financing

Figure 42 : Total expenditure on health as percentage of GDP in the African Region, 2010



Countries of the African Region without data are not included in the chart.

Figure 43 : Growth rate of total expenditure on health as percentage of GDP in the African Region, 2000 and 2010



Countries of the African Region without data are not included in the chart.

Figure 44 : Trend in total expenditure on health as percentage of GDP in the African Region, 2004 to 2009






Figure 45 : Total expenditure on health as percentage of GDP by WHO Region, 2000 and 2009





Health financing

Figure 46 : Per capita total expenditure on health (PPP int. \$) in the African Region, 2000 and 2010



Countries of the African Region without data are not included in the chart.

Figure 47 : Per capita total expenditure on health (PPP int. \$) by WHO Regions, 2000 and 2009




Figure 48 : Per capita total expenditure on health (PPP int. \$) in the African Region, 2010






Figure 49 : Trend in average per capita total expenditure on health (PPP int. \$) in the African Regions, 2004 to 2009





Health financing

Figure 50 : General government expenditure on health as percentage of total government expenditure in the African Region, 2000 and 2010



Countries of the African Region without data are not included in the chart.

Figure 51 : General government expenditure on health as percentage of total government expenditure by WHO Region, 2000 and 2009




Figure 52 : General government expenditure on health as percentage of total government expenditure in the African Region, 2010






Figure 53 : Trend in general government expenditure on health as percentage of total government expenditure in the African Region, 2004 to 2009





Health financing

Figure 54 : General government expenditure on health as percentage of total expenditure on health in the African Region, 2000 and 2010




Figure 55 : General government expenditure on health as percentage of total expenditure on health by WHO Region, 2000 and 2009




Figure 56 : General government expenditure on health as percentage of total expenditure on health in the African Region, 2010






Figure 57 : Trend in general government expenditure on health as percentage of total expenditure on health in the African Region, 2004 to 2009





Health financing

Figure 58 : External resources for health as percentage of total expenditure on health in the African Region, 2000 and 2010



Countries of the African Region without data are not included in the chart.

Figure 59 : External resources for health as percentage of total expenditure on health by WHO Region, 2000 and 2009




Figure 60 : External resources for health as percentage of total expenditure on health in the African Region, 2010







Figure 61 : Trend in external resources for health as percentage of total expenditure on health in the African Region, 2004 to 2009





Health financing

Figure 62 : Out-of-pocket expenditure as percentage of private expenditure on health in the African Region, 2000 and 2010



Countries of the African Region without data are not included in the chart.

Figure 63 : Out-of-pocket expenditure as percentage of private expenditure on health by WHO Region, 2000 and 2009




Figure 64 : Out-of-pocket expenditure as percentage of private expenditure on health in the African Region, 2010






Figure 65 : Trend in out-of-pocket expenditure as percentage of private expenditure on health in the African Region, 2004 to 2009





Health financing

Figure 66 : Private expenditure on health as percentage of total expenditure on health in the African Region, 2000 and 2010



Countries of the African Region without data are not included in the chart.

Figure 67 : Private expenditure on health as percentage of total expenditure on health by WHO Region, 2000 and 2009




Figure 68 : Private expenditure on health as percentage of total expenditure on health in the African Region, 2010






Figure 69 : Trend in private expenditure on health as percentage of total expenditure on health in the African Region, 2004 to 2009





Health financing

Figure 70 : Per capita government expenditure on health (PPP int. \$) in the African Region, 2000 and 2010



Countries of the African Region without data are not included in the chart.

Figure 71 : Per capita government expenditure on health (PPP int. \$) by WHO Region, 2000 and 2009




Figure 72 : Per capita government expenditure on health (PPP int. \$) in the African Region, 2010






Figure 73 : Trend in per capita government expenditure on health (PPP int. \$) in the African Region, 2004 to 2009




Health workforce

Figure 74 : Physician-to-population ratio (per 10,000 population) in the African Region, 2005-2010



Countries of the African Region without data are not included in the chart.

Figure 76 : Nursing and midwifery personnel-to-population ratio (per 10,000 population) in the African Region, 2005-2010



Countries of the African Region without data are not included in the chart.

Figure 75 : Physician-to-population ratio (per 10,000 population) by WHO Region, 2005-2010





Figure 77 : Nursing and midwifery personnel-to-population ratio (per 10,000 population) by WHO Region, 2005-2010






Health workforce

Figure 78 : Dentistry personnel density (per 10,000 population) in the African Region, 2005–2010



Countries of the African Region without data are not included in the chart.

Figure 80 : Pharmaceutical personnel density (per 10,000 population) in the African Region, 2005–2010



Countries of the African Region without data are not included in the chart.

Figure 79 : Dentistry personnel density (per 10,000 population) by WHO Region, 2005–2010





Figure 81 : Pharmaceutical personnel density (per 10,000 population) by WHO Region, 2005–2010



Medical products and equipment

Figure 82 : Median percentage availability of selected generic medicines in a sample of health facilities in the African Region, countries with data in 2001-2009






Figure 83 : Median consumer price ratio of selected generic medicines (ratio of median local unit price to management sciences for health international reference price), countries with data, 2001-2009





Medical products and equipment

Figure 84 : Hospital beds per 10,000 population in the African Region, 2005–2011



Countries of the African Region without data are not included in the chart.

Figure 85 : Hospital beds per 10,000 population by WHO Region, 2005–2011




Figure 86 : Psychiatric beds (per 10 000 population) in the African Region, 2005–2011





Figure 87 : Psychiatric beds (per 10 000 population) in the WHO Regions, 2005–2011



Medical products and equipment

Figure 88 : Radiotherapy units per 1,000,000 population in the African Region, 2010






Figure 90 : Percentage of civil registration coverage for births in the African Region, 2005–2010



Health information

Figure 89 : Percentage of civil registration coverage for deaths in the African Region, 2000–2009



Countries of the African Region without data are not included in the chart.

Health technology

Figure 91 : Availability of national list of approved medical devices for procurement or reimbursement, African Region, 2010




Figure 92 : Availability of technical specifications of medical devices to support procurement or donations in the African Region, 2010




Figure 93 : Presence of health technology (medical device) national policy, African Region, 2010






Figure 94 : Presence of national guidelines, policies or recommendations on the procurement of medical devices, African Region, 2010






Health technology

Figure 95 : Procurement of medical devices carried out at the national level, African Region, 2010




No Data
 None
 Yes

Figure 96 : Presence of units in ministries of Health responsible for the implementation of the health technology national policy in the African Region, 2010




Data not available
 No Data
 None
 Yes

Figure 97 : Availability of national standards or recommended lists of medical devices in the African Region, 2010



No list available
 Yes, for different healthcare facilities
 Yes, for different healthcare facilities and specific procedures
 Yes, for specific procedures
 data not available

Figure 98 : Presence of types of lists recommending health technology in the African Region, 2010



More than one list
 No list available
 Public health emergency situations
 data not available

4. Specific programmes and services



HIV/AIDS

Figure 99 : HIV/AIDS mortality rate (per 100,000 population) in the African Region, 2009

- 1.3 - 117.0
- 117.1 - 248.0
- 248.1 - 351.0
- 351.1 - 627.0
- 627.1 - 680.0
- No Data




Figure 100 : HIV/AIDS mortality rate (per 100,000 populations) by WHO Region, 2000 and 2009




Figure 101 : Prevalence of HIV (per 100,000 population) by WHO Region, 2000 and 2009





Figure 102 : Prevalence of HIV (per 100,000 population) in the African Region, 2000 and 2009



Countries of the African Region without data are not included in the chart.

Figure 103 : Percentage of people with advanced HIV infection receiving antiretroviral (ARV) combination therapy in the African Region, 2010

- 1.0 - 1.0
- 1.1 - 16.0
- 16.1 - 43.0
- 43.1 - 72.0
- 72.1 - 93.0
- >95
- No Data




Figure 106 : Percentage of people receiving antiretroviral therapy in the African Region, 2007 and 2010




Figure 104 : Percentage of antiretroviral therapy among people with HIV/AIDS in need of treatment by WHO Region, 2007 and 2010






Figure 105 : Trend in antiretroviral therapy coverage (%) among HIV-infected pregnant women for PMTCT* in the African Region, 2005 to 2010



*Prevention of mother-to-child transmission


Countries of the African Region without data are not included in the chart.

Figure 107 : Prevalence of condom use by adults aged 15–49 years during higher-risk sex in the African Region, 2005-2009



Countries of the African Region without data are not included in the chart.

Figure 108 : Population aged 15–24 years of age with comprehensive knowledge of HIV/AIDS (%) in the African Region, by sex, 2005-2009




Countries of the African Region without data are not included in the chart.

Tuberculosis


2000
2010

Figure 109 : Tuberculosis mortality rate (per 100,000 population per year) among HIV-negative people in the African Region, 2000 and 2010



2000
2010

Figure 110 : Tuberculosis incidence rate per 100,000 population per year in the African Region, 2000 and 2010





Tuberculosis

Figure 111 : Tuberculosis prevalence per 100,000 per year population in the African Region, 2010




Figure 114 : Tuberculosis prevalence per 100,000 population per year in the African Region, 2000 and 2010




Figure 112 : Tuberculosis prevalence per 100,000 population per year by WHO Region, 2000 and 2010





Figure 113 : Trend in tuberculosis prevalence per 100,000 population per year in the African Region, 2004 to 2010



Tuberculosis

Figure 115 : Case-detection rate for all forms of tuberculosis (%) in the African Region, 2010




Figure 118 : Case-detection rate for all forms of tuberculosis (%) in the African Region, 2000 and 2010




Figure 116 : Case-detection rate for all forms of tuberculosis (%) by WHO Region, 2000 and 2010





Figure 117 : Trend in percentage of tuberculosis detection rate under DOTS in the African Region, 2003 to 2007



Countries of the African Region without data are not included in the chart.


Malaria

Figure 119 : Percentage of children under 5 years of age sleeping under insecticide-treated bed nets in the African Region, 2005-2009



Countries of the African Region without data are not included in the chart.

Figure 120 : Proportion of children under 5 years of age with fever being treated with antimalarial drugs in the African Region, 2005-2010



Countries of the African Region without data are not included in the chart.

Malaria

Figure 121 : Malaria mortality rate per 100 000 population in the African Region, 2008




Figure 124 : Reported cases of malaria (in thousands) in the African Region, 2010




Figure 122 : Malaria mortality rate by WHO Region, 2006 and 2008




Figure 123 : Trend in reported clinical malaria cases (in millions) in the African Region, 1990 to 2009



Countries of the African Region without data are not included in the chart.

Immunization and vaccines development

Figure 125 : Percentage of neonates protected at birth against neonatal tetanus in the African Region, 2011




Figure 128 : Percentage of neonates protected at birth against neonatal tetanus in the African Region, 1990 and 2011




Figure 126 : Percentage of neonates protected at birth against neonatal tetanus by WHO Region, 1990 and 2011





Figure 127 : Trend in percentage of neonates protected at birth against neonatal tetanus in the African Region, 1980 to 2011



Immunization and vaccines development

Figure 129 : Percentage immunization coverage among 1-year-olds for DTP3 in the African Region, 2011




Figure 132 : Percentage immunization coverage among 1-year-olds for DTP3 in the African Region, 1990 and 2011




Figure 130: Percentage immunization coverage among 1-year-olds for DTP3 by WHO Region, 1990 and 2011





Figure 131 : Trend in percentage immunization coverage among 1-year-olds for DTP3 in the African Region, 1980 to 2011



Immunization and vaccines development

Figure 133 : Percentage immunization coverage among 1-year-olds for HepB3 in the African Region, 2011




Figure 136 : Percentage immunization coverage among 1-year-olds for HepB3 in the African Region, 2000 and 2011




Figure 134 : Percentage immunization coverage among 1-year-olds for HepB3 by WHO Region, 1990 and 2011





Figure 135 : Trend in percentage immunization coverage among 1-year-olds for HepB3 in the African Region, 1997 to 2011



Immunization and vaccines development

Figure 137 : Percentage immunization coverage among 1-year-olds for BCG in the African Region, 2011




Figure 140 : Percentage immunization coverage among 1-year-olds for BCG in the African Region, 1990 and 2011




Figure 138: Percentage immunization coverage among 1-year-olds for BCG by WHO Region, 1990 and 2011





Figure 139 : Trend in percentage immunization coverage among 1-year-olds for BCG in the African Region, 1980 to 2011



Immunization and vaccines development

Figure 141: Percentage immunization coverage among 1-year-olds for Hib in the African Region, 2010




Figure 144 : Percentage immunization coverage among 1-year-olds for Hib in the African Region, 1990 and 2011




Figure 142 : Percentage immunization coverage among 1-year-olds for Hib by WHO Region, 1990 and 2011





Figure 143 : Trend in percentage immunization coverage among 1-year-olds for Hib in the African Region, 2000 to 2011



Immunization and vaccines development

Figure 145 : Percentage immunization coverage among 1-year-olds for Measles in the African Region, 2010




Figure 148 : Percentage immunization coverage among 1-year-olds for Measles in the African Region, 1990 and 2011




Figure 146 : Percentage immunization coverage among 1-year-olds for Measles by WHO Region, 1990 and 2011





Figure 147: Trend in percentage immunization coverage among 1-year-olds for Measles in the African Region, 1980 to 2011




Countries of the African Region without data are not included in the chart.

Immunization and vaccines development

Figure 149 : Percentage immunization coverage among 1-year-olds for PCV in the African Region, 2010



Figure 152 : Percentage immunization coverage among 1-year-olds for PCV in the African Region, 1990 and 2011



Countries of the African Region without data are not included in the chart.

Figure 150: Percentage immunization coverage among 1-year-olds for PCV by WHO Region, 1990 and 2011





Figure 151: Trend in percentage immunization coverage among 1-year-olds for PCV in the African Region, 1980 to 2011



Immunization and vaccines development

Figure 153 : Percentage immunization coverage among 1-year-olds for Polio in the African Region, 2010




Figure 156 : Percentage immunization coverage among 1-year-olds for Polio in the African Region, 1990 and 2011




Figure 154: Percentage immunization coverage among 1-year-olds for Hib by WHO Region, 1990 and 2011





Figure 155 : Trend in percentage immunization coverage among 1-year-olds for Polio in the African Region, 2000 to 2011



Child and adolescent health

Figure 157 : Distribution of causes of death among children aged under 5 years (in percentage) in the African Region, 2010




Figure 158 : Percentage Infants exclusively breastfed for the first 6 months of life in the African Region, 2005-2011





Figure 159 : Percentage of infants exclusively breastfed for the first 6 months of life by WHO Region, 2005-2011





Figure 160 : Percentage of infants exclusively breastfed for the first 6 months of life in the African Region, 2005-2011



Countries of the African Region without data are not included in the chart.

Maternal and newborn health

Figure 161 : Percentage of low-birth-weight newborns by WHO Region, 2005–2010






Figure 162 : Percentage of low-birth-weight newborns in the African Region, 2005–2010



Countries of the African Region without data are not included in the chart.

Figure 163 : Malnutrition prevalence, weight for age (% of children under 5) in the African Region, 2000–2009



Countries of the African Region without data are not included in the chart.

Maternal and newborn health

Figure 164 : Percentage of births attended by skilled health personnel in the African Region, 2005-2011




Figure 167 : Percentage of births attended by skilled health personnel in the African Region, 1990-1999 and 2005-2011




Figure 165 : Percentage of births attended by skilled health personnel by WHO Region, 1990-1999 and 2005-2011





Figure 166 : Estimated workforce shortage to attain 95% skilled birth attendance by 2015 in the African Region, 2010




Countries of the African Region without data are not included in the chart.

Countries of the African Region without data are not included in the chart.


Maternal and newborn health

Figure 168 : Percentage of antenatal care coverage, at least one visit, in the African Region, 2005-2011



Countries of the African Region without data are not included in the chart.

Figure 169 : Percentage of antenatal care coverage, at least four visits, in the African Region, 2005-2011



Countries of the African Region without data are not included in the chart.

Maternal and newborn health

Figure 170 : Percentage of births by caesarean section in the African Region, 2005–2010




Figure 173 : Percentage of births by caesarean section in the African Region, 2005-2010




Figure 171 : Percentage of births by caesarean section by WHO Region, 2005–2010





Figure 172 : Distribution of main causes of maternal death in the African Region, 2004



Countries of the African Region without data are not included in the chart.

Gender and women's health

Figure 174 : Adolescent fertility rate (per 1,000 girls, 15–19 years of age) by WHO Region, 2005-2010




Figure 176 : Adolescent fertility rate (per 1,000 girls, 15–19 years of age) in the African Region, 2005-2010





Figure 175 : Total women of childbearing age (15–49 years), in thousands, in the African Region, 2010



Countries of the African Region without data are not included in the chart.

Countries of the African Region without data are not included in the chart.

Gender and women's health

Figure 177 : Contraceptive prevalence in the African Region, 2005-2010




Figure 178 : Contraceptive prevalence by WHO Region, 2005-2010






Figure 179 : Percent of contraceptive use (modern method) among married women 15-49 years old in the African Region, 2000-2009



Countries of the African Region without data are not included in the chart.

Figure 180 : Contraceptive prevalence in the African Region, 2005-2010



Countries of the African Region without data are not included in the chart.

Gender and women's health

Figure 181 : Total fertility rate (average number of children) per woman in the African Region, 2010




Figure 184 : Total fertility rate per woman in the African Region, 1990 and 2010




Figure 182 : Total fertility rate per woman by WHO Region, 1990 and 2010





Figure 183 : Trend in total fertility rate per woman in the African Region, 2004 to 2010



Neglected tropical diseases

Figure 185 : Number of reported cases of leprosy in the African Region, 2010




Figure 188 : Number of reported cases of leprosy in the African Region, 2007 and 2010




Figure 186 : Number of reported cases of leprosy by WHO Region, 2010





Figure 187 : Trend in number of reported cases of leprosy (in thousands) in the African Region, 2005 to 2010



Countries of the African Region without data are not included in the chart.

Non-communicable diseases and conditions

Figure 189 : Distribution of causes of non-communicable burden of diseases (percentage of total DALYs) in the African Region, 2004




Figure 190 : Distribution of causes of neuropsychiatric burden of diseases (percentage of total DALYs) in the African Region, 2004





Figure 191 : Distribution of causes of intentional and non-intentional injuries (percentage of total DALYs) in the African Region, 2004



Figure 192 : Insufficiently active in persons aged 15 years or older, by sex, in the African Region, 2008



Countries of the African Region without data are not included in the chart.

5. Key determinants

Risk factors for health

Figure 193 : Percentage of current tobacco use in persons 15 years of age or older by WHO Region, 2009




Figure 194 : Percentage of current tobacco use in persons 15 years of age or older in the African Region, 2009




Figure 195 : Alcohol consumption (litres per person) among adults aged 15 years of age or older by WHO Region, 2005






Figure 196 : Alcohol consumption (litres per person) among adults aged 15 years of age or older in the African Region, 2008



Countries of the African Region without data are not included in the chart.

Food safety and nutrition

Figure 197 : Percentage of underweight children under 5 years of age in the African Region, 1990-1995 and 2005-2011



Countries of the African Region without data are not included in the chart.

Figure 198 : Percentage of underweight children under 5 years of age in the African Region, 2005-2011




Figure 199 : Percentage of stunted children under 5 years of age in the African Region, 2005-2011





Figure 200 : Percentage of overweight children under 5 years of age in the African Region, 2005-2011



Demography

Figure 201 : Annual growth rate (in percent) of population in the African Region, 1989-1999 and 2000-2010




Figure 202 : Annual growth rate (in percent) of population by WHO Region, 1989-1999 and 2000-2010




Figure 203 : Percentage of population living in urban areas in the African Region, 2010





Figure 204 : Trend in percentage of population living in urban areas in the African Region, 2005 to 2010



Demography

Figure 205 : Total fertility rate per woman in the African Region, 1990 and 2010




Figure 206 : Total fertility rate per woman, by WHO Region, 1990 and 2010




Figure 207 : Total fertility rate per woman in the African Region, 2010






Figure 208 : Trend in total fertility rate per woman in the African Region, 2000 to 2009



Resources and infrastructure

Figure 209 : Gross national income per capita (PPP int. \$) in the African Region, 1990 and 2010



Countries of the African Region without data are not included in the chart.

Figure 210 : Gross national income per capita (PPP int. \$) by WHO Region, 1990 and 2010




Figure 211 : Gross national income per capita (PPP int. \$) in the African Region, 2010





Figure 212 : Trend in gross national income per capita (PPP int. \$) in the African Region, 2003 to 2010





Resources and infrastructure

Figure 213 : Electrification rate in the African Region, 2009






Figure 215 : Electrification rate in the African Region, 2000-2005



Countries of the African Region without data are not included in the chart.


Figure 214 : Paved roads as percentage of all roads in the African Region, 2000-2007



Countries of the African Region without data are not included in the chart.

Poverty and income inequality

Figure 216 : Percentage of the population living under \$1 (PPP int. \$) a day (i.e. in absolute poverty) in the African Region, 2005-2009



Countries of the African Region without data are not included in the chart.

Figure 217 : Percentage of the population living under \$1 (PPP int. \$) a day (i.e. in absolute poverty) by WHO Region, 2005-2009





Figure 218 : Share of incomes by poorest and richest section of the population in the African Region, 1989-2005




Source : World Bank 2010

Countries of the African Region without data are not included in the chart.

■ Poorest 10%
■ Poorest 20%
■ Richest 10%
■ Richest 20%


Environment

Figure 219 : Percentage of population using an improved drinking water source in the African Region, 1990 and 2010



Countries of the African Region without data are not included in the chart.

Figure 220 : Percentage of urban and rural population with access to improved drinking-water source in the African Region, 2010



Countries of the African Region without data are not included in the chart.

Environment

Figure 221 : Percentage of the population using improved drinking water sources by WHO Region, 1990 and 2008




Figure 222 : Percentage of the urban and rural population with access to improved drinking-water source by WHO Region, 2008




Figure 223 : Percentage of the population using improved sanitation facilities by WHO Region, 1990 and 2008




Figure 224 : Percentage of the population using improved sanitation facilities by WHO Region, urban and rural, 2008




Figure 225 : Trend in percentage of the urban and rural population with access to improved drinking-water source in the African Region, 2000 to 2008






Figure 226 : Trend in percentage of the population using improved sanitation facilities in the African Region, urban and rural, 2000 to 2008




Environment

Figure 227 : Percentage of the population using improved sanitation facilities in the African Region, 1990 and 2010



Countries of the African Region without data are not included in the chart.


Figure 228 : Percentage of the population using improved sanitation facilities in the African Region, urban and rural, 2010



Countries of the African Region without data are not included in the chart.


Science and technology

Figure 229 : Population who are telephone fixed subscribers (per 10,000) in the African Region, 2009



Source: World Telecommunication/ICT Indicators Data | International Telecommunications Union, 2011
Countries of the African Region without data are not included in the chart.

Figure 230 : Population who are Internet users (per 10,000) in the African Region, 2008



Source: Millennium Development Goals Database | United Nations Statistics Division, 2011
Countries of the African Region without data are not included in the chart.

Emergencies and Disasters

Figure 231 : Under-five mortality by WHO Region, both sexes, 1990, 2000 and 2009






Figure 232 : The total number of internally displaced people in the African Region, 2001 and 2009



Countries of the African Region without data are not included in the chart.

Figure 233 : The total number of refugees (in thousands) in the African Region, by country of origin, 2010



Countries of the African Region without data are not included in the chart.

6. Progress on the MDGs

Health-MDGs

MDG-4 : Reduce child mortality

1990
2010





Figure 236 : Annual average rate of reduction of under-5 mortality in the African Region, 1990 and 2010




Countries of the African Region without data are not included in the chart.

■ No progress ■ Insufficient progress ■ On track

Data Source: WHO, World Health Statistics 2011, Geneva, World Health Organization, 2011.
Country and regional assessments of progress towards MDG 4 are based on average annual rates of reduction (AARR) in USMR observed for 1990-2008 and required during 2009-2015 in order to reach the MDG target of reducing USMR by two thirds by 2015, according to the following thresholds: *On track*: USMR is less than 40, or USMR is 40 or more and AARR observed for 1990-2008 is 4.0 per cent or more. *Insufficient progress*: USMR is 40 or more and AARR observed for 1990-2008 is between 1.0 per cent and 3.9 per cent. *No progress*: USMR is 40 or more and AARR observed for 1990-2008 is less than 1.0 per cent. **Cape Verde, Algeria, Mauritius and Seychelles** are not shown in the graph, but they are *On track*, their USMR are less than 40.

2010
MDG Target 2015



Health-MDGs

MDG-5: Improve maternal health

Figure 237 : Maternal mortality ratio (per 100 000 live births) in WHO Regions in 2010





Figure 239 : Annual average rate of reduction of maternal mortality (per 100 000 live births) in the African Region, by country, 1990 and 2010




Countries of the African Region without data are not included in the chart.

■ No progress ■ Making progress
■ Insufficient progress ■ On track

Data source: Trends in Maternal mortality:1990 to 2008: Estimates developed by WHO, UNICEF, UNFPA and the World Bank, WHO, 2010 and WHO, World Health Statistics 2011, Geneva, World Health Organization 2011. Countries with MMR≥100 in 1990 are categorized as "on track or with sufficient progress" if there has been 5.5% decline or more annually, "medium progress" if MMR has declined between 2% and 5.5%, making "some progress" if MMR has declined by less than 2% annually, and having "no progress" if there has been no decline in MMR. Botswana and Mauritius with MMR<100 in 1990 are not categorized. No data available for Sao Tome and Principe and Seychelles.

■ 2010
■ MDG Target 2015

Figure 238 : Maternal mortality ratio (per 100 000 live births) in the African Region, by country, 2010 and the MDG target




Health-MDGs

MDG-5 : Improve maternal health

■ 2005-2011
■ MDG Target 2015


Figure 240 : Percentage of births attended by skilled health personnel in the African Region, by country, 2005-2011 and the MDG Target 2015



Countries of the African Region without data are not included in the chart.

■ 2005-2010
■ 1990-1999

Figure 241 : Percentage of unmet need for family planning in the African Region, by country, 1990-1999 and 2005-2010




Countries of the African Region without data are not included in the chart.

MDG -6: Combat HIV/AIDS, malaria and other diseases

■ 2005-2009
■ 2000-2004


Figure 242 : Percentage of children under 5 years of age sleeping under insecticide-treated bed nets in the African Region, 2000-2004 and 2005-2009



Countries of the African Region without data are not included in the chart.

■ 2005-2010
■ 2000-2004

Figure 243 : Percentage of children under 5 years of age with fever being treated with antimalarial drugs in the African Region, 2000-2004 and 2005-2010



Countries of the African Region without data are not included in the chart.

Figure 244 : Malaria mortality rate (per 100 000 population) in the African Region, by country, 2008





Figure 245 : Trend in death cases associated with malaria in African Region, 1990-2009 *




Source : *The data are from the World Bank 2012

Health MDGs

MDG-6: Combat HIV/AIDS, malaria and other diseases

Figure 246 : Percentage of pregnant women living with HIV infection who are receiving antiretroviral therapy for preventing mother-to-child transmission, 2010



2009
2007

Figure 247 : Percentage of people 15-49 years of age living with HIV in the African Region, by country, 2007 and 2009.





Figure 248 : Trend in contraceptive prevalence rate in African Region, 2000-2010*



Source : * Data are from the World Bank 2012

Countries of the African Region without data are not included in the chart.

Health MDGs

MDG -6: Combat HIV/AIDS, malaria and other diseases.

Figure 249 : Prevalence of tuberculosis (per 100 000 population) in the African Region, by country, 2009






Figure 251 : Trend in Tuberculosis case detection rate (% , all forms) in Sub-Saharan Africa ,1990 - 2010 *



Source : * The data are from the World Bank, 2012, unless the value of 2010 is of GHO (African Region)

Figure 250 : Percentage of population living with advanced HIV infection who are receiving antiretroviral therapy, 2010



Health-Related MDGs

MDG-7: Ensure environmental sustainability

■ 2010
■ MDG Target 2015

Figure 252 : Percentage of the population using improved drinking water sources in the African Region, by country, 2010 and the MDG target 2015





Figure 253 : Percentage of the population using improved sanitation facilities in the African Region, by country, 2010 and the MDG target 2015



Health-Related MDGs

MDG-1: Eradicate extreme poverty and hunger

■ 2005-2011
■ MDG Target 2015

Figure 254 : Percentage of underweight children under 5 years of age in the African Region, by country, 2005-2011 and the MDG target




Figure 255 : Percentage of children aged under 5 years that are underweight in the African Region, by country, 2005-2011





Figure 256 : Annual average rate of reduction of underweight children under 5 years of age in the African Region, by country, 1990-1995 and 2005-2011



Data source: WHO, World Health Statistics 2011, Geneva, World Health Organization 2011.
Countries are classified as: "On track" if average annual rate of reduction (AARR) is greater than or equal to 2.6%; "Making progress" if AARR is between 0.6% and 2.5%; "insufficient progress" if AARR is between 0.0% and 0.5% "No progress" if AARR is less than to 0.0%.

Health-Related MDGs

MDG-8: Develop a global partnership for development

Figure 257 : Official development assistance (ODA) received as percentage of GDP in the African Region, by country, 2010






Figure 259 : Trade (% of GDP) in Sub-Saharan Africa, 1990-2010*



Source : * The data are from the World Bank, 2012

1990
2009

Figure 258 : Total debt service as percentage of exports of goods, services and income in the African Region, by country, 1990 and 2009



Countries of the African Region without data are not included in the chart.

Health-Related MDGs

MDG-8: Develop a global partnership for development

Figure 260 : Percentage of the population who are Internet users in the African Region, by country, 2010




Figure 262 : Unemployment, youth total (% of total labor force ages 15-24) in the African Region, by country, 1990-1999 and 2000-2009





Figure 261 : Cellular or mobile subscribers (per 100 population) in the African Region, by country, 2010



7. Explanatory notes

The following provides the definition of the health statistics categories included in this volume, as well as the rationale for their inclusion and the estimation methods used in their production.

Health status

1. Life expectancy at birth

Rationale for use: Life expectancy at birth reflects the overall mortality level of a population. It summarizes the mortality pattern that prevails across all age groups, children and adolescents, adults and the elderly.

Definition: Average number of years that a newborn is expected to live if current mortality rates continue to apply.

Methods of estimation: WHO has developed a model life table based on about 1800 life tables from vital registration judged to be of good quality. For countries with vital registration, the level of completeness of recorded mortality data in the population is assessed and mortality rates are adjusted accordingly. Where vital registration data for 2003 were available, these were used directly to construct the life table. For countries where the information system provided a time series of annual life tables, parameters from the life table were projected using a weighted regression model, giving more weight to recent years. Projected values of the two life table parameters were then applied to the modified logit life table model, where the most recent national data provided an age pattern, to predict the full life table for 2003. In case of inadequate sources of age-specific mortality rates, the life table is derived from estimated under-5 mortality rates and adult mortality rates that are applied to a global standard (defined as the average of all the 1800 life tables using a modified logit model).

Source: Life Tables for WHO Member States. Geneva, WHO, 2010. Available at: www.who.int/whosis/database/life_tables/life_tables.cfm.

2. Healthy life expectancy (HALE)

Rationale for use: Substantial resources are devoted to reducing the incidence, duration and severity of major diseases that cause morbidity but not mortality and to reducing their impact on people's lives. It is important to capture both fatal and nonfatal health outcomes in a summary measure of average levels of population health. Healthy life expectancy (HALE) at birth adds up expectation of life for different health states, adjusted for severity distribution, making it sensitive to changes over time or differences between countries in the severity distribution of health states.

Definition: Average number of years that a person can expect to live in 'full health' by taking into account years lived in less than full health due to disease and/or injury.

Methods of estimation: Since comparable health state prevalence data are not available for all countries, a four-stage strategy is used. Data from the WHOGBD study are used to estimate severity-adjusted prevalence by age and sex for all countries. Data from the WHOMCSS and WHS are used to make independent estimates of severity-adjusted prevalence by age and sex for survey countries. Prevalence for all countries is calculated based on GBD, MCSS and WHS estimates. Life tables constructed by WHO are used with Sullivan's method to compute HALE for countries. HALE estimates use methods described in the statistical annex to the World Health Report 2004. Estimates for 2007 have been revised to take into account the Global Burden of Disease estimates for Member States for the year 2004 and may not be entirely comparable with those for 2002 published in World Health Statistics 2007. Income-group aggregates are based on the 2008 World Bank list of economies.

3. Life table (see Life expectancy at birth)

Data sources: (i) civil or sample registration: mortality by age and sex is used to calculate age-specific rates. (ii) Census: mortality by age and sex is tabulated from questions on recent deaths that occurred in the household during a given period preceding the census (usually 12 months). (iii) Census or surveys: direct or indirect methods provide adult mortality rates based on information on survival of parents or siblings.

Methods of estimation: Empirical data from different sources are consolidated to obtain estimates of the level and trend in adult mortality by fitting a curve to the observed mortality points. However, to obtain the best possible estimates, judgement needs to be made on data quality and how representative it is of the population. Recent statistics based on data availability in most countries are point estimates dated by at least 3–4 years, which need to be projected forward in order to obtain estimates of adult mortality for the current year. When no adequate source of age-specific mortality exists, the life table is derived as described in the life expectancy indicator.

4. Probability of dying (per 1000) between ages 15 and 60 years (adult mortality rate)

Rationale for use: Disease burden from noncommunicable diseases among adults – the most economically productive age span – is rapidly increasing in developing countries owing to ageing and health transitions. Therefore, the level of adult mortality is becoming an important indicator for the comprehensive assessment of the mortality pattern in a population. **Definition:** Probability that a 15-year-old person will die before reaching his/her 60th birthday. Mortality data: World Health Organization, 2010 (www.who.int/healthinfo/statistics/mortality/en/).

5. Probability of dying (per 1000) under age 5 years (under-5 mortality rate)/Probability of dying (per 1000) under age one year (infant mortality rate)

Rationale for use: Under-5 and infant mortality rates are leading indicators of the level of child health and overall development in countries. They are also MDG indicators.

Definition: The under-5 mortality rate is the probability of a child born in a specific year or period dying before reaching the age of 5, if subject to age-specific mortality rates of that period. The infant mortality rate is the probability of a child born in a specific year or period dying before reaching the age of one, if subject to age-specific mortality rates of that period.

Methods of estimation: Empirical data from different sources are consolidated to obtain estimates of the level and trend in under-5 mortality by fitting a curve to the observed mortality points. However, to obtain the best possible estimates, judgement needs to be made on data quality and how representative it is of the population. Recent statistics based on data availability in most countries are point estimates dated by at least 3–4 years, which need to be projected forward in order to obtain estimates of under-5 mortality for the current year. Those are then converted to their corresponding infant mortality rates through model life table systems: the one developed by WHO for countries with adequate vital registration data and Coale–Demeny model life tables for the other countries. It should be noted that the infant mortality data from surveys are exposed to recall bias; hence their estimates are derived from under-5 mortality, which leads to a supplementary step to estimate infant mortality rates.

6. Maternal mortality ratio (per 100 000 live births)

Rationale for use: Complications during pregnancy and childbirth are leading causes of death and disability among women of reproductive age in developing countries. The maternal mortality ratio (MMR) represents the risk associated with each pregnancy, i.e. the obstetric risk. It is also an MDG indicator for monitoring Goal 5 of improving maternal health.

Definition: Number of maternal deaths per 100 000 live births during a specified time period, usually one year.

Methods of estimation: Measuring maternal mortality accurately is difficult except where comprehensive registration of deaths and their causes exists. Elsewhere, censuses or surveys can be used to measure levels of maternal mortality. Data derived from health services records are problematic where not all births take place in health facilities, because of biases whose dimensions and direction cannot be determined. Reproductive age mortality studies (RAMOS) use triangulation of different sources of data on deaths of women of reproductive age, including record review and/or verbal autopsy, to accurately identify maternal deaths. Based on multiple sources of information, RAMOS are considered the best way to estimate levels of maternal mortality. Estimates derived from household surveys are usually based on information retrospectively collected about the deaths of sisters of the respondents and could refer back up to an average 12 years, and they are subject to wide confidence intervals. For countries without any reliable data on maternal mortality, statistical models are applied. Global and regional estimates of maternal mortality are developed every 5 years, using a regression model.

Sources: (i) Towards Reaching Health-Related Millennium Development Goals: Progress Report and Way Forward. Report of the Regional Director. Brazzaville: WHO Regional Office for Africa, 2009. (ii) Maternal Mortality in 2005. Estimates Developed by WHO, UNICEF, UNFPA and the World Bank. Geneva: WHO, 2008. Available at: http://whqlibdoc.who.int/publications/2007/9789241596213_eng.pdf.

7. Age-standardized death rates per 100 000 by cause

Rationale for use: The numbers of deaths per 100 000 population are influenced by the age distribution of the population. Two populations with the same age-specific mortality rates for a cause of death will have different overall death rates if the age distributions of their populations are different. Age-standardized mortality rates adjust for differences in population age distribution by applying the observed age-specific mortality rates for each population to a standard population.

Definition: The age-standardized mortality rate is a weighted average of the age-specific mortality rates per 100 000 persons, where the weights are the proportions of persons in the corresponding age groups of the WHO standard population. Rates are age-standardized to WHO's World Standard Population. See Age Standardization of Rates: A New WHO Standard. Geneva, WHO, 2001 (GPE Discussion Paper Series No. 31). Available at: www.who.int/healthinfo/paper31.pdf.

8. Years of life lost (percentage of total)

Rationale for use: Years of life lost (YLL) take into account the age at which deaths occur by giving greater weight to deaths at younger age and lower weight to deaths at older age. The years of life lost (percentage of total) indicator measures the YLL due to a cause as a proportion of the total YLL lost in the population due to premature mortality.

Definition: YLL are calculated from the number of deaths multiplied by a standard life expectancy at the age at which death occurs. The standard life expectancy used for YLL at each age is the same for deaths in all regions of the world and is the same as that used for the calculation of disability-adjusted life-years (DALYs). Additionally, 3% time discounting and non-uniform age weights that give less weight to years lived at young and older ages were used as for the DALY. With non-uniform age weights and 3% discounting, a death in infancy corresponds to 33 YLL, and deaths at ages 5–20 to around 36 YLL.

Source: Mortality and Burden of Disease Estimates for WHO Member States in 2004. Geneva, WHO, 2009.

Available at: www.who.int/entity/healthinfo/statistics/bodgbddeathdalyestimates.xls. Communicable diseases include maternal causes, conditions arising during the perinatal period and nutritional deficiencies. Income-group aggregates are based on the 2004 World Bank list of economies. Individual percentages may not add up to 100% owing to rounding.

9. The disability-adjusted life-year (DALY)

Rationale for use: DALY is a health gap measure that extends the concept of potential years of life lost due to premature death (PYLL) to include equivalent years of 'healthy' life lost by virtue of being in states of poor health or disability. DALYs for a disease or health condition are calculated as the sum of the years of life lost due to premature mortality (YLL) in the population and the years lost due to disability (YLD) for incident cases of the health condition.

Methods of estimation: Life tables specifying all-cause mortality rates by age and sex for 192 WHO Member States were developed for 2002 from available death registration data, sample registration systems (India and China) and data on child and adult mortality from censuses and surveys. Cause-of-death distributions were estimated from death registration data for 107 countries, together with data from population-based epidemiological studies, disease registers and notification systems for selected specific causes of death. Causes of death for populations without useable death registration data were estimated using cause-of-death models together with data from population-based epidemiological studies, disease registers and notification systems for 21 specific causes of death.

10. Causes of death among children under 5 years of age (%)

Rationale for use: MDG4 consists in the reduction of under-5 mortality by two-thirds in 2015, from its level in 1990. Child survival efforts can be effective only if they are based on reasonably accurate information about the causes of childhood deaths. Cause-of-death information is needed to prioritize interventions and plan for their delivery, to determine the effectiveness of disease-specific interventions, and to assess trends in disease burden in relation to national and international goals.

Definition: The cause(s) of death (CoD) as entered on the medical certificate of cause of death in countries with civil (vital) registration system. The underlying CoD is being analysed. In countries with incomplete or no civil registration, causes of death are those reported as such in epidemiological studies that use verbal autopsy algorithms to establish CoD.

Methods of estimation: CoD data from civil registration systems were evaluated for their completeness. Complete and nationally representative data were then grouped by ICD codes into the cause categories, and their proportions to total under-5 deaths were then computed. For countries with incomplete or no data, the distribution of deaths by cause was estimated in two steps. In the first step, a statistical model was used to assign deaths to one of three broad categories of causes: communicable diseases; non-communicable diseases; or injuries and external causes. In a second step, cause-specific under-5 mortality estimates from the Child Health Epidemiology Reference Group (CHERG), WHO Technical Programmes and the Joint United Nations Programme on HIV/AIDS (UNAIDS) were taken into account in assigning the distribution of deaths to specific causes. A variety of methods, including proportional mortality and natural history models, were used by CHERG and WHO to develop country-level cause-specific mortality estimates. All CHERG working groups developed comparable and standardized procedures to generate estimates from the databases.

Source: Mortality Data. Geneva, WHO, 2010. Available at: www.who.int/healthinfo/statistics/mortality/en/.

The health system

Health financing

11. Total expenditure on health as percentage of gross domestic product (GDP)

12. General government expenditure on health as percentage of total general government expenditure

13. Per capita total expenditure on health at international dollar rate

Rationale for use: Health financing is a critical component of health systems. National health accounts (NHAs) provide a large set of indicators based on the expenditure information collected within an internationally recognized framework. NHAs are a synthesis of the financing and spending flows recorded in the operation of a health system, from funding sources to the distribution of funds across providers and functions of health systems and benefits across geographic, demographic, socioeconomic and epidemiological dimensions.

Definitions:

- Total health expenditure as percentage of GDP
- Percentage of total general government expenditure that is spent on health
- Per capita total expenditure on health at international dollar rate

Methods of estimation: Only about 95 countries either have produced a full NHA or report expenditure on health to the Organisation for Economic Cooperation and Development (OECD). Standard accounting estimation and extrapolation techniques have been used to provide time series. The principal international references used are the International Monetary Fund (IMF) Government Finance Statistics and International Financial Statistics; OECD Health Data and International Development Statistics; and the United Nations National Accounts Statistics. National sources include national health accounts reports, public expenditure reports, statistical yearbooks and other periodicals, budgetary documents, national accounts reports, statistical data on official websites, central bank reports, non-governmental organization reports, academic studies, and reports and data provided by central statistical offices and ministries.

Source: WHO National Health Accounts (NHA), Country Health Expenditure Database. Geneva: WHO, February 2010.

Available at: www.who.int/nha/country/. The regional, income and global figures are calculated using Purchasing Power Parity (PPP) terms. When the number is smaller than 0.05%, the percentage may appear as zero. For per capita expenditure indicators, this is represented as <1. In countries where the fiscal year begins in July, expenditure data have been allocated to the later calendar year (for example, 2008 data will cover the fiscal year 2007–08). Absolute values of expenditures are expressed in nominal terms (current prices). National currency units per US\$ are calculated using the average exchange rates for the year. For 2008, the use of yearly average exchange rates (compared with year-end exchange rates) may not fully represent the impact of the global financial crisis.

14. General government expenditure on health as percentage of total expenditure on health

15. General government expenditure on health as percentage of total government expenditure

16. External resources for health as percentage of total expenditure on health

17. Out-of-pocket expenditure as percentage of private expenditure on health

18. Per capita total expenditure on health at average exchange rate (US\$)

19. Per capita government expenditure on health at average exchange rate (US\$)

20. Per capita government expenditure on health at international dollar rate

Rationale for use: Health financing is a critical component of health systems. NHAs provide a large set of indicators based on the expenditure information collected within an internationally recognized framework. NHAs are a synthesis of the financing and spending flows recorded in the operation of a health system, from funding sources to the distribution of funds across providers and functions of health systems and benefits across geographic, demographic, socioeconomic and epidemiological dimensions.

Definitions: Key indicators for which the data are available:

- Level of total expenditure on health as percentage of GDP, and per capita health expenditures in US dollars and in international dollars.
- Distribution of public and private sectors in financing health and their main components, such as:

*Extent of social and private health insurance

*Burden on households through out-of-pocket spending

*Reliance on external resources in financing health

Associated terms:

-Gross domestic product (GDP) is the value of all goods and services provided in a country by residents and non-residents. This corresponds to the total sum of expenditure (consumption and investment) of the private and government agents of the economy during the reference year.

-General government expenditure (GGE) includes consolidated direct outlays and indirect outlays, such as subsidies and transfers, including capital, of all levels of government social security institutions, autonomous bodies, and other extrabudgetary funds.

-Total expenditure on health (THE) is the sum of general government health expenditure and private health expenditure in a given year, calculated in national currency units in current prices. It comprises the outlays earmarked for health maintenance or for restoration or enhancement of the health status of the population, paid for in cash or in kind

-General government expenditure on health (GGHE) is the sum of outlays by government entities to purchase health care services and goods. It comprises the outlays on health by all levels of government and by social security agencies, and direct expenditure by parastatals and public firms. Expenditures on health include final consumption, subsidies to producers and transfers to households (chiefly reimbursements for medical and pharmaceutical bills). It includes both recurrent and investment expenditures (including capital transfers) made during the year. Besides domestic funds, it also includes external resources (mainly as grants passing through the government or loans channelled through the national budget).

-Social security expenditure on health (SSHE) includes outlays for purchases of health goods and services by schemes that are mandatory and controlled by government. Such social security schemes that apply only to a selected group of the population, such as public sector employees only, are also included here.

-External resources health expenditure (ExtHE) includes all grants and loans, whether passing through governments or private entities for health goods and services, in cash or in kind.

-Private health expenditure (PvtHE) is defined as the sum of expenditures on health by the following entities:

*Prepaid plans and risk-pooling arrangements (PrepaidHE) are the outlays of private insurance schemes and private social insurance schemes (with no government control over payment rates and participating providers, but with broad guidelines from government)

*Firms' expenditure on health are the outlays by private enterprises for medical care and health-enhancing benefits other than payment to social security or other prepaid schemes.

*Non-profit institutions serving mainly households are the outlays of those entities whose status do not permit them to be a source of financial gain for the units that establish, control or finance them. This includes funding from internal and external sources.

*Household out-of-pocket spending (OOPS) comprises the direct outlays of households, including gratuities and in-kind payments made to health practitioners and to suppliers of pharmaceuticals, therapeutic appliances, and other goods and services. This includes household direct payments to public and private providers of health care services, nonprofit institutions, and non-reimbursable cost sharing, such as deductibles, copayments and fee for services.

- Exchange rate is the annual average or year-end number of units at which a currency is traded in the banking system.

- International dollars are derived by dividing local currency units by an estimate of their Purchasing Power Parity (PPP) compared with the US dollar, i.e. the measure that minimizes the consequences of differences in price levels between countries.

Methods of estimation and sources: About 100 countries either have produced full national health accounts or report expenditure on health to the OECD. Standard accounting estimation and extrapolation techniques have been used to provide time series (1998–2004). Ministries of Health have responded to the draft updates sent for their inputs and comments. For details on sources and methods, see www.who.int/nha.

Health workforce

21. Number of:

- physicians per 10000 population

- nurses per 10000 population

- midwives per 10000 population

Rationale for use: The availability and composition of human resources for health is an important indicator of the strength of the health system. Even though there is no consensus about the optimal level of health workers for a population, there is ample evidence that worker numbers and quality are positively associated with immunization coverage, outreach of primary care, and infant, child and maternal survival.

Definitions:

- Physicians: includes generalists and specialists.

- Nurses: includes professional nurses, auxiliary nurses, enrolled nurses and other nurses, such as dental nurses and primary care nurses.

- Midwives: includes professional midwives, auxiliary midwives and enrolled midwives. Traditional birth attendants, who are counted as community health workers, appear elsewhere.

- Dentists: includes dentists, dental assistants and dental technicians.

- Pharmacists: includes pharmacists, pharmaceutical assistants and pharmaceutical technicians.

- Public and environmental health workers: includes environmental and public health officers, sanitarians, hygienists, public and environmental health technicians, district health officers, malariatechnicians, meat inspectors, public health supervisors, and similar professions.

- Community health workers: includes traditional medicine practitioners, faith healers, assistant/community health education workers, community health officers, family health workers, lady health visitors, health extension package workers, community midwives, institution-based personal care workers and traditional birth attendants.

- Laboratory health workers: includes laboratory scientists, laboratory assistants, laboratory technicians and radiographers.

- Other health workers: includes a large number of occupations such as dieticians and nutritionists, medical assistants, occupational therapists, operators of medical and dentistry equipment, optometrists and opticians, physiotherapists, podiatrists, prosthetic/orthotic engineers, psychologists, respiratory therapists, speech pathologists, and medical trainees and interns.

- Health management and support workers: includes general managers, statisticians, lawyers, accountants, medical secretaries, gardeners, computer technicians, ambulance staff, cleaning staff, building and engineering staff, skilled administrative staff, and general support staff.

Methods of estimation: No methods of estimation have been developed.

Source: WHO Global Atlas of the Health Workforce. Geneva: WHO, 2009. Available at: <http://apps.who.int/globalatlas/default.asp>. See this source for the latest updates, time-trend statistics and disaggregated data, as well as metadata descriptors. In general, the denominator data for health workforce density (i.e. national population estimates) were obtained from the World Population Prospects Database of the United Nations Population Division. In some cases, official reports provided only workforce density indicators, from which estimates of the absolute numbers were calculated. Depending on the organization of national health systems and means of monitoring, data may not be exactly comparable across countries. Data from the years prior to 2000 were excluded from this edition.

Medical Products and Equipment

22. Number of hospital beds per 10 000 population

Rationale for use: Service delivery is an important component of health systems. To capture availability access and distribution of health services delivery, a range of indicators or a composite indicator is needed. Currently, there are no such data for the majority of countries. Inpatient bed density is one of the few available indicators on a component of level of health service delivery.

Definition: Number of inpatient beds per 10 000 population.

Methods of estimation: Empirical data only, with possible adjustment for underreporting (e.g. missing private facilities). Additional data are compiled by the WHO Regional Office for Africa.

Health Information – Civil Registration Coverage

23. Coverage of vital registration of deaths

Rationale for use: Health information is an essential component of health systems. The registration of births and deaths with causes of death, called 'civil registration (vital registration)', is an important component of a country's health information system.

Definition: Percentage of estimated total deaths that are 'counted' through a civil registration system.

Methods of estimation: Expected numbers of deaths by age and sex are estimated from current life tables, based on multiple sources. Reported numbers are compared with expected numbers by age and sex to obtain an estimate of coverage of the vital registration system.

Sources: (i) **United Nations Demographic Yearbook 2007**. New York: United Nations Statistics Division, 2009.

Available at: <http://unstats.un.org/unsd/demographic/products/dyb/dybsets/2007%20DYB.pdf>; (ii) WHO Mortality Database: Tables. Geneva: WHO, 2009.

Available at: www.who.int/healthinfo/mortables.

Specific programmes and services

24. Estimated rate of adults (15 years and older) dying of HIV/AIDS (per 1000)/Estimated rate of children below 15 years of age dying of HIV/AIDS (per 1000)

Rationale for use: The mortality rates for adults and children below 15 are leading indicators of the level of impact of the HIV/AIDS epidemic and the impact of interventions, especially scale-up of treatment and prevention of mother-to-child transmission in countries.

Definition: Estimated mortality due to HIV/AIDS is the number of adults and children that have died in a specific year based in the modelling of HIV surveillance data using standard and appropriate tools.

Methods of estimation: Empirical data from different HIV surveillance sources are consolidated to obtain estimates of the level and trend in adult and child mortality by using standard methods and tools for HIV estimates appropriate to the level of HIV epidemic. However, to obtain the best possible estimates, judgement needs to be made on data quality and how representative it is of the population. UNAIDS/WHO produce country-specific estimates every 2 years.

Source: Based on the 2008 Report on the Global AIDS Epidemic. Geneva, UNAIDS and WHO, 2008. See Annex: HIV and AIDS Estimates and Data, 2007 and 2001. Available at: http://data.unaids.org/pub/GlobalReport/2008/jc1510_2008_global_report_pp211_234_en.pdf. Ranges of estimates are available from this document. WHO regional and global figures are updates for the year 2008. Income-group aggregates are based on the 2008 World Bank list of economies.

25. HIV prevalence among the population aged 15–49 years

Rationale for use: HIV/AIDS has become a major public health problem in many countries, and monitoring the course of the epidemic and the impact of interventions is crucial. Both the MDGs and the United Nations General Assembly Special Session on HIV and AIDS (UNGAS) have set goals for reducing HIV prevalence.

Definition: Percentage of people with HIV infection among all people aged 15–49 years.

Methods of estimation: HIV prevalence data from HIV sentinel surveillance systems, which may include national population surveys with HIV testing, are used to estimate HIV prevalence using standardized tools and methods of estimation developed by UNAIDS and WHO in collaboration with the UNAIDS Reference Group on Estimation, Modelling and Projections. Tools for estimating the level of HIV infection are different for generalized epidemics, and concentrated or low-level epidemics.

26. People with advanced HIV infection receiving antiretroviral (ARV) combination therapy (%)

Rationale for use: As the HIV epidemic matures, increasing numbers of people are reaching advanced stages of HIV infection. ARV combination therapy has been shown to reduce mortality among those infected, and efforts are being made to make it more affordable even in less-developed countries. This indicator assesses the progress in providing ARV combination therapy to everyone with advanced HIV infection.

Definition: Percentage of people with advanced HIV infection receiving ARV therapy according to nationally approved treatment protocol (or WHO/Joint UN Programme on HIV and AIDS standards) among the estimated number of people with advanced HIV infection.

Methods of estimation: The denominator of the coverage estimate is obtained from models that also generate the HIV prevalence, incidence and mortality estimates. The number of adults with advanced HIV infection who need to start treatment is estimated as the number of AIDS cases in the current year times 2. The total number of adults needing ARV therapy is calculated by adding the number of adults who need to start ARV therapy to the number of adults who are being treated in the previous year and have survived into the current year.

Source: Towards Universal Access: Scaling Up Priority HIV/AIDS Interventions in The Health Sector: Progress Report, 2008. Geneva: WHO, Joint United Nations Programme on HIV/AIDS, United Nations Children's Fund, 2008. WHO regional and global figures are updates for the year 2008. Income-group aggregates are based on the World Bank 2008 list of economies.

27. Incidence of tuberculosis

Rationale for use: Incidence (cases arising in a given time period) gives an indication of the burden of TB in a population, and of the size of the task faced by a national TB control programme. Incidence can change as the result of changes in transmission (the rate at which people become infected with *Mycobacterium tuberculosis*, the bacterium that causes TB) or changes in the rate at which people infected with *M. tuberculosis* develop TB disease (e.g. as a result of changes in nutritional status or of HIV infection). Because TB can develop in people who became infected many years previously, the effect of TB control on incidence is less immediate than the effect on prevalence or mortality. MDG6, Target 8 is 'to have halted by 2015 and begun to reverse the incidence of TB. WHO estimates that in 2004 the per capita incidence of TB was stable or falling in 5 out of 6 WHO Regions, but growing globally at 0.6% per year. The exception was the African Region, where incidence is apparently still increasing, but less rapidly each year. Implementation of the Stop TB Strategy, following the Global Plan to Stop TB 2006–2015, is expected to reverse the rise in incidence globally by 2015.

Definition: Estimated number of TB cases arising in a given time period (expressed as per capita rate). All forms of TB are included, as are cases in people with HIV.

Methods of estimation: Estimates of TB incidence, prevalence and mortality are based on a consultative and analytical process in WHO and are published annually. Estimates of incidence for each country are derived using one or more of four approaches, depending on the available data:

incidence = case notifications/proportion of cases detected

incidence = prevalence/duration of condition

incidence = annual risk of TB infection × Stýblo coefficient

incidence = deaths/proportion of incident cases that die

Data are for all forms of tuberculosis, including tuberculosis in people with HIV infection.

Source: Global Tuberculosis Control: A Short Update to the 2009 Report. Geneva: WHO, 2009 (WHO/HTM/TB/2009.426).

Available at: www.who.int/tb/publications/global_report. WHO region, income group and global aggregates include territories.

28. Prevalence of tuberculosis

Rationale for use: Prevalence and mortality are direct indicators of the burden of TB, indicating the number of people suffering from the disease at a given point in time and the number dying each year. Furthermore, prevalence and mortality respond quickly to improvements in control, as timely and effective treatments reduce the average duration of disease (thus decreasing prevalence) and the likelihood of dying from the disease (thus reducing disease-specific mortality). MDG6 is 'to combat HIV/AIDS, malaria and other diseases' (including TB). This goal is linked to Target 8, 'to have halted by 2015 and begun to reverse the incidence of malaria and other major diseases', and MDG Indicator 24, 'prevalence and mortality rates associated with TB'. The Stop TB Partnership has endorsed the related targets of reducing per capita TB prevalence and mortality by 50% relative to 1990, by the year 2015. There are few good data with which to establish TB prevalence and mortality, particularly

for the baseline year of 1990. However, current best estimates suggest that implementation of the Global Plan to Stop TB 2006–2015 will halve 1990 prevalence and mortality rates globally and in most regions by 2015, though not in Africa and Eastern Europe.

Definition: The number of cases of TB (all forms) in a population at a given point in time (sometimes referred to as 'point prevalence') expressed in this database as number of cases per 100 000 population.

Methods of estimation: Estimates of TB incidence, prevalence and mortality are based on a consultative and analytical process in WHO and are published annually. The methods used to estimate TB prevalence and mortality rates are described in detail elsewhere. Country-specific estimates of prevalence are, in most instances, derived from estimates of incidence, combined with assumptions about the duration of disease. The duration of disease is assumed to vary according to whether the disease is smear-positive or not; whether the individual receives treatment in a DOTS programme or non-DOTS programmes, or is not treated at all; and whether the individual is infected with HIV.

29. Tuberculosis: DOTS case detection rate

Rationale for use: The proportion of estimated new smear-positive cases that are detected (diagnosed and notified to WHO) by DOTS programmes provides an indication of how effective national TB programmes are in finding people with TB and diagnosing the disease.

Methods of estimation: Estimates of incidence are based on a consultative and analytical process in WHO, and are published annually. The DOTS detection rate for new smear-positive cases is calculated by dividing the number of new smear-positive cases notified to WHO by the estimated number of incident smear-positive cases for the same year.

30. Tuberculosis: DOTS treatment success

Rationale for use: Treatment success is an indicator of the performance of national TB control programmes. In addition to the obvious benefit to individual patients, successful treatment of infectious cases of TB is essential to prevent the spread of the infection. Detecting and successfully treating a large proportion of TB cases should have an immediate impact on TB prevalence and mortality. By reducing transmission, successfully treating the majority of cases will also affect, with some delay, the incidence of disease.

Definition: The proportion of new smear-positive TB cases registered under DOTS in a given year that successfully completed treatment, whether with bacteriological evidence of success ('cured') or without ('treatment completed'). At the end of treatment, each patient is assigned one of the following six mutually exclusive treatment outcomes: cured; completed; died; failed; defaulted; and transferred out with outcome unknown. The proportions of cases assigned to these outcomes, plus any additional cases registered for treatment but not assigned to an outcome, add up to 100% of cases registered. The treatment-success rate is the percentage of new smear-positive patients registered for treatment who were cured (with laboratory confirmation) or who completed their course of treatment.

Source: Global TB Control: A Short Update to the 2009 Report. Geneva: WHO, 2009 (WHO/HTM/TB/2009.426).

Available at: www.who.int/tb/publications/global_report. WHO Regional, income group and global aggregates include territories.

31. Tuberculosis mortality

Rationale for use: Prevalence and mortality are direct indicators of the burden of tuberculosis (TB), indicating the number of people suffering from the disease at a given point in time and the number dying each year. Furthermore, prevalence and mortality respond quickly to improvements in control, as timely and effective treatment reduces the average duration of disease (thus decreasing prevalence) and the likelihood of dying from the disease (thus reducing disease-specific mortality).

Definition: Estimated number of deaths due to TB in a given time period. It is expressed in this database as deaths per 100 000 population per year. Includes deaths from all forms of TB, and deaths from TB in people with HIV.

Methods of estimation: Estimates of TB incidence, prevalence and mortality are based on a consultative and analytical process in WHO and are published annually. The methods used to estimate TB mortality rates are described in detail elsewhere. Country-specific estimates of TB mortality are, in most instances, derived from estimates of incidence, combined with assumptions about the case fatality rate. The case fatality rate is assumed to vary according to whether the disease is smear-positive or not; whether the individual receives treatment in a DOTS programme or non-DOTS programmes, or is not treated at all; and whether the individual is infected with HIV. These are classified as deaths from tuberculosis according to the International Statistical Classification of Diseases and Related Health Problems, 10th revision. Geneva, WHO, 1992.

Source: Global Tuberculosis Control: A Short Update to the 2009 Report. Geneva, WHO, 2009 (WHO/HTM/TB/2009.426).

Available at: www.who.int/tb/publications/global_report. WHO Regional, income group and global aggregates include territories.

32. Children under 5 years of age with fever who received treatment with any antimalarial therapy (%)

Rationale for use: Prompt treatment with effective antimalaria drugs for children with fever in malaria risk areas is a key intervention to reduce mortality. In addition to being listed as a global MDG indicator under Goal 6, effective malaria treatment is also identified by WHO, UNICEF and the World Bank as one of the four main interventions to reduce the burden of malaria in Africa: (i) use of insecticide-treated nets (ITNs); (ii) prompt access to effective treatments in or near the home, (iii) provision of antimalaria drugs to symptom-free pregnant women in stable transmission areas; and (iv) improved forecasting, prevention and response, essential to respond quickly and effectively to malaria epidemics. In areas of sub-Saharan Africa with stable levels of malaria transmission, it is essential that access to prompt treatment be ensured. This requires drug availability at household or community level and, for complicated cases, availability of transport to the nearest equipped facility. Reserve drug stocks, transport and hospital capacity are needed to mount an appropriate response to malaria cases and prevent the onset of malaria from degenerating to a highly lethal complicated malaria picture.

Definition: Percentage of population under 5 years of age in malaria-risk areas with fever being treated with effective antimalaria drugs.

Methods of estimation: For prevention, the indicator is calculated as the percentage of children under 5 years of age who received effective antimalaria drugs upon a fever episode. The information is obtained directly from household surveys. The empirical values are directly reported without further estimation.

33. Children under 5 years of age sleeping under insecticide-treated nets (%)

Rationale for use: In areas of intense malaria transmission, malaria-related morbidity and mortality are concentrated in young children, and the use of insecticide-treated nets (ITNs) by children under 5 years of age has been demonstrated to considerably reduce malaria disease incidence, malaria-related anaemia and all-cause under-5 mortality. Vector control through the use of ITNs constitutes one of the four intervention strategies of the Roll Back Malaria Initiative. It is also listed as an MDG indicator.

Definition: Percentage of children under 5 years of age in malaria-endemic areas who slept under an ITN the previous night, ITN being defined as a mosquito net that has been treated within 12 months or is a long-lasting insecticidal net (LLIN).

Methods of estimation: Empirical data only.

Source: World Malaria Report 2009, Annex 6. Geneva: WHO, 2009.

Available at: www.who.int/malaria/world_malaria_report_2009/mal2009_annex6_0010.pdf.

34. Number of poliomyelitis cases

Rationale for use: the 1988 World Health Assembly (WHA) called for the global eradication of poliomyelitis. The number of poliomyelitis cases is used to monitor progress towards this goal and to inform eradication strategies. Countries implement strategies supplementing routine immunization (e.g. national

immunization days and sub-national campaigns) or more targeted mop-up activities, depending on the levels of poliomyelitis cases.

Definition: Suspected polio cases (acute flaccid paralysis (AFP), other paralytic diseases, and contacts with polio cases) that are confirmed by laboratory examination or are consistent with polio infection.

Methods of estimation: Estimates of polio cases are based exclusively on unadjusted surveillance data.

Source: Data from WHO Polio Eradication Initiative, as of 12 January 2010.

Updated information can be found at: www.who.int/immunization_monitoring/en/diseases/poliomyelitis/case_count.cfm. Confirmed polio cases refer to any circulating polioviruses (wild poliovirus and circulating vaccine-derived poliovirus (cVDPV)).

35. One-year-olds immunized with:

- one dose of measles (%)

- three doses of diphtheria, tetanus toxoid and pertussis (DTP3) (%)

- three doses of hepatitis B (HepB3)(%)

Rationale for use: Immunization coverage estimates are used to monitor immunization services and to guide disease eradication and elimination efforts, and are a good indicator of health systems performance.

Definition: Measles immunization coverage is the percentage of 1-year-olds who have received at least one dose of measles containing vaccine in a given year. For countries recommending the first dose of measles among children older than 12 months of age, the indicator is calculated as the proportion of children less than 24 months of age receiving one dose of measles containing vaccine. DTP3 immunization coverage is the percentage of 1-year-olds who have received three doses of the combined diphtheria and tetanus toxoid and pertussis vaccine in a given year. HepB3 immunization coverage is the percentage of 1-year-olds who have received three doses of Hepatitis B3 vaccine in a given year.

Methods of estimation: WHO and UNICEF rely on reports from countries, household surveys and other sources such as research studies. Both organizations have developed common review process and estimation methodologies. Draft estimates are made, reviewed by country and external experts and then finalized.

Sources: Unless otherwise stated, data are derived from Demographic and Health Surveys (DHS) conducted since 2000. The DHS figures were extracted using STATA compiler software (www.measuredhs.com/). When not available using STATA compiler software, figures were extracted directly from DHS reports. For some countries and some of the indicators, there were differences in the figures extracted from the country reports and STATA compiler. In these cases, following discussions with staff from the MEASURE DHS implementation group (ICF Macro), data from the country reports were used. Further information regarding the source of individual country data can be obtained on request from WHO.

36. Antenatal care coverage (%)

Rationale for use: Antenatal care coverage is an indicator of access and utilization of health care during pregnancy.

Definition: Percentage of women who utilized antenatal care provided by skilled health personnel for reasons related to pregnancy at least once during pregnancy as a percentage of live births in a given time period.

Methods of estimation: Empirical data from household surveys are used. At global level, facility data are not used.

Source: UNICEF Global Database on Maternal Health. New York: UNICEF, 2010. Available at: www.childinfo.org/antenatal_care_country.php.

37. Births by caesarean section (%)

Rationale for use: The proportion of births by caesarean section is an indicator of access to and utilization of health care during childbirth.

Definition: Percentage of births by caesarean section among all live births in a given time period.

Methods of estimation: Empirical data from household surveys.

38. Births attended by skilled health personnel (%)

Rationale for use: All women should have access to skilled care during pregnancy and at delivery to ensure detection and management of complications. Moreover, because it is difficult to measure maternal mortality accurately, model-based maternal mortality ratio (MMR) estimates cannot be used for monitoring short-term trends. The proportion of births attended by skilled health personnel is used as a proxy indicator for this purpose.

Definition: Percentage of live births attended by skilled health personnel in a given period of time.

Methods of estimation: Empirical data from household surveys are used. At a global level, facility data are not used.

Source: WHO Global Database on Maternal Health Indicators, 2009 update. Geneva: WHO, 2009.

Available at: www.who.int/reproductivehealth/global_monitoring/index.html. In order to enhance comparability over time, the reported figures are derived, to the extent possible, from broadly comparable data sources. Therefore, reported figures may not refer to the most recently available data. Refer to the source for more complete information on time trends and metadata.

39. Contraceptive prevalence (%)

Rationale for use: Contraceptive prevalence is an indicator of health, population, development and women's empowerment. It also serves as a proxy measure of access to reproductive health services that are essential for meeting many of the MDGs, especially the child mortality, maternal health, HIV/AIDS and gender-related goals.

Definition: Contraceptive prevalence is the proportion of women of reproductive age who are using (or whose partner is using) a contraceptive method at a given point in time

Methods of estimation: Empirical data only.

Source: World Contraceptive Use 2009. New York: Population Division, Department of Economic and Social Affairs, United Nations Secretariat, 2009 (POP/DB/CP/Rev2009).

40. Condom use at higher-risk sex among young people aged 15–24 years (%)

Rationale for use: Consistent correct use of condoms within non-regular sexual partnerships substantially reduces the risk of sexual HIV transmission.

Definition: Percentage of young people aged 15–24 years reporting the use of a condom during the last sexual intercourse with a non-regular partner among those who had sex with a non-regular partner in the last 12 months.

Methods of estimation: Empirical data only. Survey respondents aged 15–24 years are asked whether they have commenced sexual activity. Those who report sexual activity and have had sexual intercourse with a non-regular partner in the last 12 months are further asked about the number of non-regular

partners and condom use the last time they had sex with a nonregular partner.

Source: Data are from Demographic and Health Surveys (DHS) and exclude country-reported data. 2008 Report on the Global AIDS Epidemic. Geneva: Joint United Nations Programme on HIV/AIDS, WHO, 2008. Available at: www.unaids.org/en/KnowledgeCentre/HIVData/GlobalReport/2008/2008_Global_report.asp. See Annex 2: Country Progress Indicators.

41. Children under 5 years of age with acute respiratory infection and fever (ARI) taken to facility

Rationale for use: Respiratory infections are responsible for almost 20% of all under-5 deaths worldwide. The number of under-5s with ARI who are taken to an appropriate health provider is a key indicator for both coverage of intervention and careseeking and provides critical inputs to the monitoring of progress towards the child-survival-related MDGs and strategies.

Definition: Proportion of children aged 0–59 months who had presumed pneumonia (ARI) in the last 2 weeks and were taken to an appropriate health provider.

Methods of estimation: Empirical data.

42. Children under 5 years of age with diarrhoea who received ORT

Rationale for use: Diarrhoeal diseases remain one of the major causes of under-5 mortality, accounting for 1.8 million child deaths worldwide, despite all the progress in their management and the undeniable success of oral rehydration therapy (ORT). Therefore, the monitoring of the coverage of this very cost-effective intervention is crucial for the monitoring of progress towards the child-survival-related MDGs and strategies.

Definition: Proportion of children aged 0–59 months who had diarrhoea in the last 2 weeks and were treated with oral rehydration salts or an appropriate household solution (ORT).

Methods of estimation: Empirical data.

43. Children 6–59 months of age who received vitamin A supplementation

Rationale for use: Vitamin A supplementation is considered a critically important intervention for child survival owing to the strong evidence that exists of its impact on child mortality. Therefore, measuring the proportion of children who have received vitamin A in the last 6 months is crucial for monitoring coverage of interventions towards the child-survival-related MDGs and strategies.

Definition: proportion of children 6–59 months of age who have received a high-dose vitamin A supplement in the last 6 months.

Methods of estimation: Empirical data.

Source: Data compiled by WHO from Demographic and Health Surveys (DHS) and Multiple Indicator Cluster Surveys (MICS), January 2010. Available at: www.measuredhs.com and www.unicef.org/statistics/index_24302.html.

44. Prevalence of current tobacco use in adolescents (13–15 years of age)

Rationale for use: The risk of chronic diseases starts early in childhood, and tobacco use continues to adulthood. Tobacco is an addictive substance, and smoking often starts in adolescence, before the development of risk perception. By the time the risk to health is recognized, the addicted individuals find it difficult to stop tobacco use.

Definition: Prevalence of tobacco use (including smoking, oral tobacco and snuff) on more than one occasion in the 30 days preceding the survey, among adolescents 13–15 years old.

Source: WHO/CDC Global Youth Tobacco Survey (GYTS). Geneva: WHO, 2010. Available at: www.cdc.gov/tobacco/global/GYTS/results.htm. Data relate to tobacco use in any form in the past 30 days.

45. Prevalence of current (daily or occasional) tobacco smoking among adults (15 years and older) (%)

Rationale for use: Prevalence of current tobacco smoking among adults is an important measure of the health and economic burden of tobacco, and provides a baseline for evaluating the effectiveness of tobacco control programmes over time. While a more general measure of tobacco use, including both smoked and smokeless products, would be ideal, data limitations restrict the present indicator to smoked tobacco. Occasional tobacco smoking constitutes a significant risk factor for tobacco-related disease, and is therefore included along with daily tobacco smoking.

Definition: Prevalence of current tobacco smoking (including cigarettes, cigars, pipes or any other smoked tobacco products). Current smoking includes both daily and non-daily or occasional smoking.

Methods of estimation: Empirical data only.

Source: Based on WHO Report On The Global Tobacco Epidemic, 2009: Implementing Smoke-Free Environments. Geneva: WHO, 2009. Available at: www.who.int/tobacco/mpower/en/. See Appendix VII: Age-Standardized Prevalence Estimates for WHO Member States, 2006. 'Smoking' is defined as smoking at the time of the survey of any form of tobacco, including cigarettes, cigars, pipes, bidis, etc. and excluding smokeless tobacco. These figures represent age-standardized prevalence rates for smoking tobacco, and should only be used to draw comparisons of prevalence between countries and between men and women within a country. They should not be used to calculate the number of smokers in a country, region, income group or globally.

46. Children under 5 years of age

- stunted for age (%)

- underweight for age (%)

- overweight for age (%)

Rationale for use: All three indicators measure growth in young children. Child growth is internationally recognized as an important public health indicator for monitoring nutritional status and health in populations. In addition, children who suffer from growth retardation as a result of poor diets and/or recurrent infections tend to have greater risks of illness and death.

Definition: Percentage of children stunted describes how many children under 5 years have a height-for-age below minus two standard deviations of the National Center for Health Statistics (NCHS)/WHO reference median. Percentage of children underweight describes how many children under 5 years have a weight-for-age below minus two standard deviations of the NCHS/WHO reference median. Percentage of children overweight describes how many children under 5 years have a weight-for-height above two standard deviations of the NCHS/WHO reference median.

Methods of estimation: Empirical values. Several countries have limited data for recent years and current estimations are made using models that make projections based on past trends.

Source: Global Database on Child Growth and Malnutrition. Geneva: WHO, 2009. Available at: www.who.int/nutgrowthdb/database/en. Prevalence estimates are based on WHO standards.

47. Newborns with low birthweight (%)

Rationale for use: the low-birthweight rate at the population level is an indicator of a public health problem that includes long-term maternal malnutrition, ill-health and poor health care. On an individual basis, low birthweight is an important predictor of newborn health and survival.

Definition: Percentage of live-born infants with birthweight less than 2500 g in a given time period. Low birthweight may be subdivided into very low birthweight (less than 1500 g) and extremely low birthweight (less than 1000 g).

Methods of estimation: Where reliable health service statistics with a high level of coverage exist, percentage of low-birthweight births. For household survey data, different adjustments are made according to the type of information available (numerical birthweight data or subjective assessment by the mother).

Source: UNICEF Global Database on Low Birthweight. New York: UNICEF, 2009. Available at: www.childinfo.org/low_birthweight_table.php (November 2009 update).

48. Prevalence of adults (15 years and older) who are obese (%)

Rationale for use: The prevalence of overweight and obesity in adults has been increasing globally. Obese adults (BMI ≥ 30.0 kg/m²) are at increased risk of adverse metabolic outcomes, including increased blood pressure, cholesterol, triglycerides and insulin resistance. Subsequently, an increase in BMI exponentially increases the risk of noncommunicable diseases (NCDs), such as coronary heart disease, ischaemic stroke and type 2 diabetes mellitus. Raised BMI is also associated with an increased risk of cancer.

Definition: Percentage of adults classified as obese (BMI ≥ 30.0 kg/m²) among total adult population (15 years and older).

Methods of estimation: Estimates are still under development and will be published later in 2006. Only nationally representative surveys with either anthropometric data collection or self-reported weight and height (mostly in high income countries) are included in the 2006 World Health Statistics. Comparisons between countries may be limited owing to differences in sample characteristics or survey years.

Source: Global Database on Body Mass Index. Geneva, WHO, 2010. Available at: www.who.int/bmi.

Key determinants of health

49. Population with:

- sustainable access to an improved water source (%)

- access to improved sanitation (%)

Rationale for use: Access to drinking water and improved sanitation is a fundamental need and a human right vital for the dignity and health of all people. The health and economic benefits of improved water supply to households and individuals (especially children) are well documented. Both indicators are used to monitor progress towards the MDGs.

Definition: Access to an improved water source is the percentage of the population with access to an improved drinking water source in a given year. Access to improved sanitation is the percentage of the population with access to improved sanitation in a given year.

Methods of estimation: Estimates are generated through analysis of survey data and linear regression of data points. Coverage estimates are updated every 2 years.

Source: WHO/UNICEF Joint Monitoring Programme for Water Supply and Sanitation. Geneva: WHO and UNICEF, 2010. Available at: www.wssinfo.org/en/welcome.html.

50. Population using solid fuels (%)

Rationale for use: The use of solid fuels in households is associated with increased mortality from pneumonia and other acute lower respiratory diseases among children, as well as increased mortality from chronic obstructive pulmonary disease and lung cancer (where coal is used) among adults. It is also an MDG indicator.

Definition: Percentage of population using solid fuels.

Methods of estimation: The data from surveys and censuses are used as reported in the surveys and censuses. All countries with a Gross National Income (GNI) per capita above US\$ 10 500 are assumed to have made a complete transition to cooking with non-solid fuels. For low- and middle-income countries with a GNI per capita below US\$ 10 500 and for which no household solid fuel use data are available, a regression model based on GNI, percentage of rural population, and location or non-location within the Eastern Mediterranean Region is used to estimate the indicator. These estimates use methods developed and implemented by the WHO/UNICEF Joint Monitoring Programme for Water Supply and Sanitation. Where solid fuel use information is available for two or more separate years (spaced at least 5 years apart) linear regression is performed. The linear regression line is extrapolated up to 2 years after the latest survey point and up to 2 years before the earliest survey point. Outside these time limits, the extrapolated regression line is flat for 4 years in either direction. Where coverage reaches 0% or 100%, a horizontal line is drawn from the year before coverage reaches 0% or 100%. For countries with solid fuel use at less than 5%, 0% is assumed for the calculation of regional or global aggregates; for countries with more than 95%, 95% is assumed in the calculation of the aggregate.

Source: WHO Household Energy Database. Geneva: WHO, 2010. Available at: www.who.int/indoorair/health_impacts/he_database/en/.



Basic data and statistics are at the core of all health systems. Without them, it would be impossible to analyse evidence and extract action-orientated knowledge for decision making.

The development of an African Health Observatory and national health observatories aim to narrow the knowledge gap and strengthen health systems in the African Region by providing easy access to high quality information, evidence and knowledge, as well as facilitate their use for policy and decision making.

WHO Health Situation Analysis in the Africa Region: Atlas of Health Statistics, 2012 presents in numerical and graphical formats the best data available for key health indicators in the 46 countries of WHO's African Region.

With the continued input and collaboration of the African countries, this publication and its future editions will be a significant, constantly updated information product of the Observatory.