

EVERY NEWBORN

An Action Plan To End Preventable Deaths



About Every Newborn

The *Every Newborn* action plan is based on the latest epidemiology, evidence and global and country learning, and supports the United Nations Secretary-General's *Every Woman Every Child* movement. The preparation was guided by the advice of experts and partners, led by WHO and UNICEF, and by the outcome of several multistakeholder consultations and a web-based consultation with more than 300 comments. Discussed at the 67th World Health Assembly, Member States endorsed the document and made firm commitments to put in practice recommended actions. The Director General has been requested to monitor progress towards the achievement of the global goal and targets and report periodically to the Health Assembly until 2030.

WHO Library Cataloguing-in-Publication Data

Every Newborn: an action plan to end preventable deaths.

1.Infant Mortality – prevention and control. 2.Infant, Newborn. 3.Stillbirth. 4.Perinatal Care. 5.Pregnancy Outcome. I.World Health Organization

ISBN 978 92 4 150744 8 (NLM classification: WA 900)

© World Health Organization 2014

All rights reserved. Publications of the World Health Organization are available on the WHO website (www.who.int) or can be purchased from WHO Press, World Health Organization, 20 Avenue Appia, 1211 Geneva 27, Switzerland (tel.: +41 22 791 3264; fax: +41 22 791 4857; e-mail: bookorders@who.int).

Requests for permission to reproduce or translate WHO publications –whether for sale or for non-commercial distribution– should be addressed to WHO Press through the WHO website (www.who.int/about/licensing/copyright_form/en/index.html).

The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted lines on maps represent approximate border lines for which there may not yet be full agreement.

The mention of specific companies or of certain manufacturers' products does not imply that they are endorsed or recommended by the World Health Organization in preference to others of a similar nature that are not mentioned. Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters.

All reasonable precautions have been taken by the World Health Organization to verify the information contained in this publication. However, the published material is being distributed without warranty of any kind, either expressed or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall the World Health Organization be liable for damages arising from its use.

Reprinted in Luxembourg

EVERY NEWBORN An Action Plan To End Preventable Deaths

JUNE 2014

Contents

i. Foreword	3
ii. Acronyms and key definitions	
1. Executive summary	5
2. Today's landscape Existing commitments Rights Current situation	9 10
3. Effective interventions for improving the health of newborns across the continuum of care]∠
4. Vision and goals	18
5. Strategic objectives and principles Strategic objective 1 Strategic objective 2 Strategic objective 3 Strategic objective 4 Strategic objective 5	20 22 28 32
6. Framework for success Impact framework Coverage targets Measures of success Milestones Core indicators	39 40 41
7. The way forward. Research priorities. Coordination.	45
8. Actions by constituency	46
References	48
Summary of action plan process	50
ANNEX	
Annex 1: Sixty-Seventh World Health Assembly Resolution WHA67.10: Newborn health action plan	





Foreword by the United Nations Secretary-General Ban Ki-moon

The world can be proud of progress achieved since the launch of the Global Strategy for Women's and Children's Health in 2010. The *Every Woman Every Child* initiative has generated new attention and investment to address some of the most neglected causes of women's and children's mortality.

At the same time, we must confront the tragic fact that each year, 289 000 women still die while giving life and an estimated 18 000 children die every day from preventable diseases and circumstances. This is an appalling tragedy that demands an accelerated response.

It is time to give newborns a more prominent place on the global health agenda. We must do much more to save the 2.9 million newborns dying during their first 28 days of life each year. The day of birth is the most dangerous day, when nearly half of maternal and newborn deaths and stillbirths occur. It is also the day babies face the greatest risk of disability.

Protecting newborns means ensuring proper care for their mothers before, during and after pregnancy. The majority of these deaths are preventable, caused by complications related to prematurity, birth and severe infections. Preterm and small babies are in much greater peril, including the long-term risk of stunting and developing non-communicable diseases like diabetes and hypertension as adults.

The *Every Newborn* action plan takes forward the Global Strategy for Women's and Children's Health. By focusing on the quality of care at birth, we can save millions of lives and accelerate progress towards the Millennium Development Goals.

I ask governments, civil society, donors, the private sector and other partners to take action to address preventable newborn deaths and ensure that the rights of women and children are fully recognized. The strong accountability mechanism under the *Every Woman Every Child* umbrella will ensure that resources and results are tracked when it comes to commitments to newborns.

Let us do all we can to ensure a healthy start for all mothers and newborns. This will open the way for progress across the development agenda and around the world.

Acronyms

MDG Millennium Development Goal

NMR Neonatal mortality rate

RMNCAH Reproductive, maternal, newborn, child and adolescent health

SBR Stillbirth rate

WHO World Health Organization

Key definitions

Low birth weight Weight of less than 2 500 g, irrespective of gestational age

Newborn death The death within 28 days of birth of any live-born baby regardless

of weight or gestational age

Preterm birth A baby born < 37 completed weeks gestation

Small for gestational age Infant below the 10th percentile of birth weight for gestational age.

An SGA baby may be preterm or full-term.

Stillbirth A baby born with no signs of life, weighing more than 1 000 g or with more than

28 completed weeks of gestation (for international comparison purposes)





Executive summary

The promise and potential of Every Newborn: an action plan to end preventable deaths

Although remarkable progress has been made in recent decades to reduce the number of child deaths worldwide, too many newborns continue to die each year despite the availability of feasible, evidence-based solutions.

Newborn survival and health and prevention of stillbirths were not specifically addressed in the Millennium Development Goal (MDG) framework and consequently received less attention and investment. Newborn deaths and stillbirths are reducing at a slower rate than under-5 deaths and maternal deaths (1).

Now is the time for the global health community to prioritize this unfinished agenda.

Today, we have unprecedented opportunities to turn the tide and address newborn health, as far more is known about effective interventions, service delivery channels and approaches to accelerate coverage and quality of care. Recently, renewed commitments to saving newborn lives and preventing stillbirths have

been made by many governments and partners in response to the United Nations Secretary-General's Global Strategy for Women's and Children's Health (2) and its accompanying Every Woman Every Child initiative, Committing to Child Survival: A Promise Renewed (3), and to recommendations made by the Commission on Information and Accountability for Women's and Children's Health (4) and the United Nations Commission on Life-Saving Commodities for Women and Children (5). The much-needed attention specifically for newborn health has triggered multiple stakeholders to propose this Every Newborn: an action plan to end preventable deaths. The Every Newborn action plan is based on epidemiology, evidence, and global and country learning, setting a framework to end preventable newborn deaths and stillbirths by 2035.

The final plan was endorsed at the 67th World Health Assembly in May 2014 (Annex 1).

The action plan sets out a vision of a world in which there are no preventable deaths of newborns or stillbirths, where every pregnancy is wanted, every birth celebrated and women, babies and children survive, thrive and reach their full potential. Nearly 3 million lives could be saved each year if the actions in the plan are implemented and its goals and targets achieved. Based on evidence of what works, and developed within the framework for Every Woman Every Child, the plan enhances and supports coordinated, comprehensive

planning and implementation of newborn-specific actions within the context of national reproductive, maternal, newborn, child and adolescent health (RMNCAH) strategies and action plans, and in collaboration with stakeholders from the private sector, civil society, professional associations and others. The goal is to achieve equitable and high-quality coverage of care for all women and newborns through links with other global and national plans, measurement and accountability.

Key messages

3 million babies and women could be saved each year through investing in quality care around the time of birth and special care for sick and small newborns. Cost-effective solutions are now available to protect women and children from the most dangerous day of their lives – the day of birth.

Unfinished agenda: Newborn health and stillbirths are part of the "unfinished agenda" of the Millennium Development Goals for women's and children's health. With newborn deaths still accounting for 44% of under-5 deaths globally, newborn mortality and stillbirths require greater visibility in the emerging post-2015 sustainable development agenda if the overall under-5 mortality is to be reduced.

We have solutions to address the main causes of newborn death: More than 80% of all newborn deaths result from three preventable and treatable conditions – complications due to prematurity, intrapartum-related deaths (including birth asphyxia) and neonatal infections. Costeffective, proven interventions exist to prevent and treat each main cause. Improving quality of care around the time of birth will save the most lives, but this requires educated and equipped health workers, including those with midwifery skills, and availability of essential commodities.

Women's and children's health is a smart investment, particularly with specific attention to care at birth: High coverage of care around the time of birth and care of small and sick newborns would save nearly 3 million lives (women, newborns and stillbirths) each year at an additional running cost of only US\$ 1.15 per person in 75 high burden countries. This would have a triple impact on investments – saving women and newborns and preventing stillbirths.

Action with a plan: The *Every Newborn* action plan was developed in response to country demand. It sets out a clear vision of how to improve newborn health and prevent stillbirths by 2035. The plan builds on the United Nations Secretary General's Global Strategy for Women's and Children's Health and the *Every Woman Every Child* movement by supporting government leadership and providing guidance on how to strengthen newborn health components in existing health sector plans and strategies, especially those that relate to reproductive, maternal and child health. *Every Newborn* calls upon all stakeholders to take specific actions to improve access to, and quality of, health care for women and newborns within the continuum of care.

Vision, goals, guiding principles and strategic objectives

A world in which there are no preventable deaths of newborns or stillbirths, where every pregnancy is wanted, every birth celebrated, and women, babies and children survive, thrive and reach their full potential.

Goals

Goal 1: Ending preventable newborn deaths Goal 2: Ending preventable stillbirths

By 2035, all countries will reach the target of 10 or less newborn deaths per 1 000 live births 10 or less stillbirths per 1 000 total births and and continue to reduce death and disability, ensuring that no newborn is left behind.

By 2035, all countries will reach the target of continue to close equity gaps.

Guiding principles

- Country leadership
- Human rights
- Integration

- Equity
- Accountability
- Innovation

Strategic objectives

- 1. Strengthen and invest in care during labour, birth and the first day and week of life.
- 2. Improve the quality of maternal and newborn care.
- 3. Reach every woman and newborn to reduce inequities.
- 4. Harness the power of parents, families and communities.
- 5. Count every newborn through measurement, programme-tracking and accountability.

Effective interventions

We now have unprecedented opportunities for improving newborn health after decades of research that have generated solid evidence on the burden and causes of neonatal mortality, demonstrated effective interventions and service delivery channels, and identified ways to accelerate progress and scale up interventions to save lives. High-impact, cost-effective interventions for newborn health, like breastfeeding support and kangaroo mother care, form one component of integrated health services for RMNCAH.

Interventions for both woman and baby delivered at the same time in the same place

by the same health care provider/team will have the highest impact on saving lives and improving health outcomes. High coverage of interventions before, during and after pregnancy could save nearly 3 million women, stillbirths and newborns by 2025 in 75 highburden countries at an additional cost of only US\$ 1.15 per capita (6). The packages of care with greatest impact on ending preventable neonatal deaths and stillbirths include: care during labour, childbirth and the first week of life; and care for the small and sick newborn. These packages are the focus of the Every Newborn action plan as they would not only save the most newborn lives but also prevent maternal deaths and stillbirths.

Framework for success

The Every Newborn action plan establishes specific global and national targets and milestones for quality of care, newborn mortality and stillbirth rates, monitoring,

investments and the implementation of national plans to support RMNCAH. Achieving these milestones will help ensure that the vision and goals for 2035 are achieved.


Call to action

The intrinsic link between the survival and health **We must act now.** We know the main of newborns, stillbirths and the survival, health and nutrition of all women of reproductive age, including before, during, between and after pregnancy is clear. The action plan emphasizes the need to reach every woman and newborn when they are most vulnerable - during labour, birth and in the first days of life. Investment in this critical time period provides the greatest potential for ending preventable neonatal deaths, stillbirths and maternal deaths, and would result in a triple return on investment.

Health research in the delivery, development and discovery of appropriate interventions must be at the forefront of efforts to reduce newborn mortality and stillbirths. Research and innovation can help improve delivery of health services and discover new solutions to prevent preterm birth and other causes of maternal and newborn death. Improving delivery of known interventions is a top priority.

causes of newborn deaths, and we have the knowledge and tools to prevent them. It is not too late to accelerate progress towards the MDGs. We must also prioritize quality care at the time of birth for women and newborns in the post-2015 sustainable development agenda and include indicators for newborn mortality and stillbirths. A healthy society is one in which women and adolescent girls, newborns and children survive and thrive.

If we commit to working together as a global community to take the specific actions outlined in the Every Newborn action plan, we can achieve our vision of a world in which there are no preventable deaths of newborns or stillbirths, where every pregnancy is wanted, every birth celebrated, and women, babies and children survive, thrive and reach their full potential and in doing so create equitable societies and transform human development.





Today's landscape

The number of child deaths worldwide has declined markedly in recent decades, largely through interventions to lower mortality after the first month of life. The mortality rate among children under-5 years of age has fallen globally by almost 50% (from 90 deaths per 1 000 live births in 1990 to 48 deaths per 1 000 live births in 2012), but the neonatal mortality rate decreased only 37% (from 33 deaths per 1 000 live births to 21 deaths per 1 000 live births) over the same period and represented, in 2012, 44% of the total child mortality (7).

Many newborn deaths are preventable and could be avoided if the actions in this plan were implemented and its goals and targets achieved. The draft action plan is based on evidence of what works. It recognizes that survival of a newborn is a sensitive marker of a health system's response to its most vulnerable citizens and calls upon all stakeholders to improve access to, and quality of, health care for women and newborns within the continuum of care that spans pre-conception, pregnancy, childhood and adolescence.

The action plan sets out a vision of a world in which there are no preventable deaths of newborns or stillbirths, every birth celebrated, and women, babies and children survive, thrive and reach their full potential. To realize this vision, the plan proposes strategic objectives, goals and mortality targets by 2035, with intermediate targets for 2020, 2025 and 2030. Targets are also proposed for coverage of quality care for women and babies around the time of birth, care of sick and small newborns and postnatal care. The action plan includes an impact framework, milestones and indicators to measure progress.

Developed within the framework for the Every Woman Every Child initiative, the plan aims to enhance and support coordinated, comprehensive planning and implementation of newborn-specific actions within the context of national RMNCAH strategies and action plans. The goal is to achieve equitable and high-quality coverage of essential, referral and emergency care for women and newborns in every country through links with other global and national plans, measurement and accountability.

The focus is primarily on newborn survival and health and the prevention of stillbirths. These targets were not included in the MDG framework and consequently received less attention and investment, resulting in slower reductions in mortality.

Stillbirths, newborn survival and health are intrinsically linked with the survival, health and nutrition of women before conception and during and between pregnancies. The periods of greatest risk for morbidity and mortality for woman and child are the hours that precede and the hours and days that follow birth. The action plan therefore emphasizes the need to reach every woman and newborn when they are most vulnerable - in labour, during birth and in the first days of life. Intervention in this critical time period provides the greatest potential for ending preventable neonatal deaths, stillbirths and maternal deaths, and would result in a triple return on investment. The plan is part of the broader initiative to end preventable maternal and newborn deaths; strategic objectives and targets to achieve the goal of ending preventable maternal deaths have also been prepared (Annex 2). The objectives are complementary to those of the Every Newborn action plan and intended for coordinated implementation.

The action plan has been drafted in close consultation with stakeholders; a draft version uploaded on the World Health Organization (WHO) website in February 2014 elicited more than 300 formal comments. The plan takes into account all inputs, the findings from an analysis of obstacles to scaling up effective interventions to improve newborn health (8),

a comprehensive epidemiological analysis (9) and a review of the evidence of the effectiveness of the proposed interventions (6). The final plan was endorsed at the 67th World Health Assembly (WHA) in May 2014.

Existing commitments

Putting the action plan into practice will strengthen existing commitments, such as the pledge made in the Family Planning 2020 initiative (10), "A Promise Renewed" (which calls for an end to preventable child deaths) (3) and strong regional and global undertakings to ending preventable maternal deaths. Given the large proportion of deaths of under-5-year olds that occur in the neonatal period and the close link between maternal and neonatal mortality, these commitments will not be fulfilled without specific efforts to reduce neonatal mortality and stillbirths. The plan builds on, and links to, other global action plans, such as those on nutrition, vaccines, malaria, pneumonia, diarrhoea, water and sanitation, and elimination of mother-tochild transmission of HIV, syphilis and neonatal tetanus, and takes into consideration multiple timeframes for achievement of these existing commitments' targets and goals.

Many governments and partners have renewed their commitments to saving the lives of women and newborns in response to the United Nations Secretary-General's Global Strategy for Women's and Children's Health (launched in 2010) (2) and accompanying initiative Every Woman Every Child, and to the recommendations from the Commission on Information and Accountability for Women's and Children's Health (4) and the United Nations Commission on Life-Saving Commodities for Women and Children (5). Recognizing that progress in newborn health has lagged behind advances in maternal and child health, the action plan takes forward the Global Strategy's mission, supporting the call by the World Health Assembly in resolution WHA58.31, Working towards universal coverage of maternal, newborn and child health interventions (11), to commit resources and to accelerate national action to build a seamless continuum of care for RMNCAH. These initiatives also affirm

We need to reach every woman and newborn when they are most vulnerable – during labour, birth and in the first days of life

women's and children's health as matters of fundamental human rights.


The Global Investment Framework for Women's and Children's Health estimates that additional investment of US\$ 5 per person per year on RMNCAH across the continuum of care would produce a nine-fold return on investment in social and economic benefits in the highest-burden countries (12).

The Sixty-fourth World Health Assembly in 2011, in resolution WHA64.12 on WHO's role in the follow-up to the United Nations High-level Plenary Meeting of the General Assembly on the MDGs (13), expressed deep concern at the slow and uneven progress towards MDGs 4 and 5. In resolution WHA64.13 on working towards the reduction of perinatal and neonatal mortality (14) the Health Assembly observed that movement towards reducing perinatal and neonatal mortality had stagnated and requested the Director-General to promote targeted plans to increase access to high-quality and safe health services that prevent and treat perinatal and neonatal conditions. The action plan responds to that request and also reflects the Health Assembly's decisions in resolution WHA64.9 on sustainable financing structures and universal coverage (15).

Rights

Under Articles 6 and 24 of the Convention on the Rights of the Child (16), every newborn child has the inherent right to life, survival and development, the highest attainable standard of health and access to health care services for treatment and rehabilitation. The notion of legal obligations is reinforced in General Comment No. 15 by the Committee on the Rights of the Child on the child's right to the highest attainable standard of health and health care, which specifies that States have an obligation to reduce child mortality and that particular attention should be paid to neonatal mortality.

The United Nations Human Rights Council welcomed the technical guidance on the application of a human-rights based approach to the implementation of policies and programmes to reduce preventable maternal morbidity and mortality in September 2012 (17), reinforcing women's rights to access quality, appropriate and affordable services and support across the continuum of care. The Council's resolution A/HRC/22/L.27/Rev.1 on the right of the child to the enjoyment of the highest attainable standard of health affirmed the importance of applying a human-rights-



based approach to eliminating preventable maternal and child mortality and morbidity and requested Member States to renew their political commitment and take action to address the main causes of mortality and morbidity.

Current situation


In addition to 2.9 million babies who die in the first month of life (7), it is estimated that 2.6 million babies are stillborn (die in the last three months of pregnancy or during childbirth) (9) and some 289 000 women die each year from complications of pregnancy and childbirth (18). The global annual average rate of reduction in neonatal mortality since 1990 has been 2.0%, lower than that of maternal mortality (2.6%) and under-5-year old mortality (2.9%) (1).

Most newborn deaths occur in low- and middle-income countries. Two thirds of

all neonatal mortality is reported from 12 countries, six of which are in sub-Saharan Africa^a. Countries with a rate of 30 or more deaths per 1 000 live births account for 60% of all newborn deaths.

Many countries with a high burden of newborn deaths have experienced recent conflict or humanitarian emergencies. Others have weak health systems with limited infrastructure and low density of skilled health workers and the population faces high out-of-pocket expenditures. Inequitable access to quality health services for women and children results in stark disparities in mortality rates and intervention coverage between and within countries. Nevertheless, 11 low- and lower-middle-income countries have reduced their neonatal mortality rate (NMR) by more than 40% since 2000b, showing that it is possible to make rapid progress.

Fig. 1 Causes of deaths in children under 5 years of age, 2012



More than 80% of newborn deaths are in small babies (preterm or small for gestational age) in the highest burden settings.

Source: WHO Global Health Observatory, 2014 (19). Estimates are rounded, and therefore may not sum to 100%.

^a In descending order of annual number of newborn deaths: India, Nigeria, Pakistan, China, Democratic Republic of the Congo, Ethiopia, Bangladesh, Indonesia, Angola, Kenya, United Republic of Tanzania, Afghanistan.

^b Bangladesh, Cambodia, Democratic People's Republic of Korea, Egypt, El Salvador, Malawi, Mongolia, Rwanda, Senegal, Sri Lanka and United Republic of Tanzania.

Social determinants are an important factor in the health of women and newborns. Poverty, inequality and societal unrest undermine maternal and newborn care in numerous ways, such as poor nutritional status of girls and women (including during pregnancy) and inadequate housing and sanitation. Complex humanitarian emergencies cause dramatic movements of people (including pregnant women and newborns) and compromise access to functional health systems. Low education levels, gender discrimination and a lack of empowerment prevent women from seeking health care and making the best choices for their own and their children's health, resulting in perilous delays and unnecessary deaths. This action plan acknowledges the inherent links between contextual factors and maternal and newborn health and focuses primarily on health system solutions.

Three causes accounted for more than 80% of neonatal mortality in 2012 (Fig. 1): complications of prematurity, intrapartum-related neonatal deaths (including birth asphyxia) and neonatal infections (sepsis, meningitis, pneumonia, and diarrhoea) (19). Complications of prematurity are also the second leading cause of all deaths of children aged under 5 years. Annually, 15 million babies are born prematurely (20) and 32.4 million with a weight below the tenth percentile for their gestational age (21); 10 million do not breathe at birth, of which 6 million require basic neonatal resuscitation (bag and mask ventilation) (22).

The highest risks of death in utero, in the neonatal period and throughout infancy and early childhood, are faced by small and low-birth-weight babies, that is, those who are born preterm or small for gestational age, or both. More than 80% of all newborn deaths occur among small babies in southern Asia and sub-Saharan Africa (9).

Disabilities remain an important issue. Many could be prevented by adequate care during labour and childbirth and in the neonatal period. Preterm babies who survive the first month of life face higher risks of post-neonatal mortality, long-term neurodevelopmental impairment, stunting and noncommunicable disease (23). Babies who are small for gestational age face

risks of stunting and adult-onset metabolic conditions. Four million term or near-term neonates have other life-threatening conditions, including intrapartum-related brain injury, severe bacterial infection and pathological jaundice, which result in life-long impairments (9). More than a million neonates survive each year with long-term neurodevelopmental impairment. It is therefore important to look beyond survival and provide appropriate follow-up care for children affected by these conditions to ensure early detection and appropriate care and rehabilitation (9). As Article 23 of the Convention on the Rights of the Child (16) stresses, it is also important to meet the needs of mentally or physically disabled children.

Some 10% of the entire global burden of disease is related to neonatal and congenital conditions (9). As the Health Assembly recognized in resolution WHA63.17 on birth defects (24), factors leading to birth defects and congenital abnormalities can affect health outcomes for babies who survive the neonatal period, as do many of the diverse causes and determinants of congenital disorders. Prevention of birth defects and provision of care for affected children need to be integrated into existing maternal, reproductive and child health services, with social welfare provision for all who need it.



Effective interventions for improving the health of newborns across the continuum of care

Unprecedented opportunities for improving newborn health now exist after decades of analysis and research that have generated information on the burden and causes of neonatal mortality, demonstrated effective interventions and service delivery channels, and identified ways to accelerate progress in extending the coverage of interventions to reduce mortality (25).

Effective interventions for improving survival and health of newborns form one component of integrated health services for RMNCAH (Fig. 2). These are well documented across the life course and have been packaged

for levels of service delivery (26). Many are delivered from common platforms for health care delivery; integrated planning and delivery can ensure efficient and effective health services for women and children.

Focus of the Every Newborn action plan The time around birth results in the majority of maternal and newborn deaths and stillbirths as well as human capital loss. These packages have the highest impact yet some of the lowest coverage of equitable and quality care across the continuum. Skilled care at birth Hospital care of childhood illness Comprehensive emergency obstetric and newborn care Skilled care at birth Prevention and management of childhood illness Basic emergency obstetric and newborn care Postnatal visits Care of small and sick newborns Home birth with skilled inselling and birth Ongoing care for the child at home care and clean NTERSECTORAL: Improved living and working conditions, including housing, water and sanitation, and nutrition; education impowerment, especially of girls; folic acid fortification; safe and healthy work environments for women and pregnant wo LABOUR, BIRTH AND LABOUR PREGNANCY CHILD **POSTNATAL** FIRST WEEK AND BIRTH **3 MILLION LIVES SAVED EACH YEAR**


Fig. 2 Packages in the continuum of care

Source: The Lancet Every Newborn Series, Mason E et al. Lancet, 2014 (27).

The packages of care with greatest impact on ending preventable neonatal deaths and stillbirths include: care during labour, around birth and the first week of life; and care for the small and sick newborn. High coverage of interventions before conception, as well as before, during and after pregnancy could

save nearly 3 million women, stillbirths and newborns by 2025 in 75 high-burden countries at an additional cost of US\$ 1.15 per capita (Fig. 3) (6). Since these packages would save the most newborn lives as well as prevent maternal deaths and stillbirths, these are the focus of the *Every Newborn* action plan.

Fig. 3 Lives that could be saved by 2025 with universal coverage of care



Source: The Lancet Every Newborn Series, Bhutta Z et al. Lancet, 2014 (6).

Interventions in the package of "care during labour, around birth and in the first week after birth" include (but are not limited to) skilled care at birth, basic and comprehensive obstetric care, management of preterm births (including the use of antenatal corticosteroids), and essential newborn care (hygienic care, thermal control, support for breastfeeding and, if required, newborn resuscitation). It is important that the interventions for mother and newborn are seen as a functional unit delivered in a narrow time window by the same health care provider (or team) and in the same place, with referral for management of complications including mother and baby together. These packages could prevent more than 1.9 million maternal and newborn deaths and stillbirths by 2025 with universal access (6).

The package of "care of small and sick newborns" comprises interventions to deal with complications arising from preterm birth and/or small for gestational age, and neonatal infections (sepsis, meningitis, pneumonia and those causing diarrhoea). Appropriate management of small and sick newborns includes extra thermal care and support for feeding for small or preterm babies, including kangaroo mother care, antibiotic treatment for infections and full supportive facility care. The last encapsulates additional feeding support (including cup and nasogastric tube feeding and intravenous fluids), infection prevention and management, safe oxygen therapy, case management of jaundice and possibly surfactant and respiratory support. Focusing on small or sick newborns could prevent almost 600 000 newborn deaths by 2025 (6).

Postnatal care provides the delivery platform for care of the normal newborn, including the promotion of healthy practices and detection of problems requiring additional care. It is provided in a different time window, often by different providers in different places. Care of the normal newborn includes early initiation of (exclusive) breastfeeding, prevention of

hypothermia, clean postnatal care practices and appropriate cord care. Close observation for 24 hours and at least three additional postnatal contacts (on day 3 (48–72 hours), between days 7 and 14, and at six weeks after birth) is recommended for all mothers and newborns to establish good care-giving practices and detect any life-threatening conditions (28).

Other interventions across the continuum of care are also vital for the survival and health of women and their babies. For example, antenatal care provides an opportunity for integrated service delivery for pregnant women, including obstetric services, but also covers infections, such as preventing, detecting and treating malaria and syphilis in pregnancy, caring for women with tuberculosis and HIV infection, preventing mother-to-child transmission of HIV and reducing harmful lifestyle practices such as smoking and alcohol use. According to WHO's World Health Statistics (29), the coverage rate of some interventions, such as vaccination with tetanus toxoid, is already high in many settings (82% of all newborns are now protected at birth against neonatal tetanus); there is therefore less potential for averting deaths (see Fig. 3), but coverage rates must remain high. Some 81% of women receive antenatal care at least once during pregnancy, but only 55% receive the recommended minimum of four visits or more (29), and the quality of care is often suboptimal.

Care before and between pregnancies affects the survival and health of women and their babies. Family planning is a vital contributor through delaying, spacing and limiting births, all of which can reduce newborn mortality and boost the health of mothers, their babies and their other children. Access to family planning and the right to control if, when and how frequently to become pregnant empower women and girls and improves babies' health and survival. Investments in family planning will contribute significantly to an overall reduction in

Care around the time of birth saves mothers and their newborns and prevents stillbirths and disability

Care for women before and during pregnancy, childbirth and the postnatal period and between pregnancies and for the newborn is best provided by a dedicated health professional qualified in midwifery. Care should be respectful and should optimize normal biological processes. In most cases, the person providing care will be a professional midwife, who will need support from a team composed of other health professionals, such as nurses, obstetricians and paediatricians, when complications arise. The team could provide all aspects of care in settings where there are no professional midwives: for these settings, the term "midwifery personnel" is used.

maternal and neonatal mortality: reducing the number of unintended pregnancies could avert 60% of maternal and 57% of child deaths (12).

Additional components of care before and between pregnancy that affect newborn health include life-skills education, nutrition, prevention and management of harmful practices (including smoking and alcohol use), identification and treatment of conditions such as sexually transmitted infections and mental illness, and tackling intimate partner violence. The nutritional status of women is of particular importance, as a woman undernourished before pregnancy is more likely to give birth to babies who are preterm or small for gestational age at birth, or both. This risk is determined partly by undernutrition of a woman in her own first 1 000 days and during her adolescence. It is essential to break the intergenerational cycle of ill health and undernutrition, especially given increasing evidence about links between low birth weight and undernutrition in the first 1 000 days of life and the rise of noncommunicable diseases in adulthood.

Preventing early unintended pregnancy in adolescent girls is a major component of efforts to improve newborn health (30). Very young mothers and their babies face greater risks from pregnancy and birth (31). Delaying

pregnancy in adolescent girls is a powerful means of saving maternal and newborn lives and empowering girls to finish their education.

Community health workers, especially in rural areas, can play an important role in bridging the gap between health services and families, and home visits made by them during pregnancy and in the first week after childbirth have been shown to have a positive impact on newborn care practices and NMR (32,33). They are also effective in detecting and referring mothers with postpartum complications and offering family-planning counselling. Other community agents, such as traditional leaders, influential family members and traditional birth attendants, also influence the demand for, and access to, skilled care.

The provision of a continuum of care throughout the life course requires seamless, functional coordination between levels of health services and the public and private sectors. Delivering health care to women and newborns requires coordination between technical programmes and initiatives and collaboration among all concerned stakeholders: governments, professional associations, civil society, academic and research institutions, the business community, development partners and families.



Vision and goals


The vision of the *Every Newborn* action plan is of a world in which there are no preventable deaths of newborns or stillbirths, where every pregnancy is wanted, every birth celebrated, and women, babies and children survive, thrive and reach their full potential.

Goal 1: ending preventable newborn deaths

All countries will reach the target NMR of 10 or less newborn deaths per 1 000 live births by 2035 and continue to reduce death and disability, ensuring that no newborn is left behind. Achievement of this target will result in an average global NMR of 7 deaths per 1 000 live births, a figure that is consistent with, and necessary to the achievement of, the target set in *Committing to Child Survival: A Promise Renewed* of ending preventable child deaths. By 2030, all countries will reach 12 or less

newborn deaths per 1 000 live births resulting in an average global NMR of 9 deaths per 1 000 live births (other interim targets are shown in Fig. 4). It is intended that these targets will also link to the proposed goals for ending preventable maternal deaths (Annex 2). All countries should ensure this goal is also achieved for underserved populations, maximizing human capital. The goal may be considered a continuation of MDG 4 to cover the unfinished business of reducing newborn deaths.

Fig. 4 Ending preventable newborn and child deaths




Source: The Lancet Every Newborn Series, Lawn JE et al. Lancet, 2014 (9).

Goal 2: ending preventable stillbirths

All countries will reach the target stillbirth rate (SBR) of 10 or less stillbirths per 1 000 total births by 2035 and continue to close equity gaps. Achieving this goal will result in an average global SBR of 8 per 1 000 total births. By 2030, all countries will reach 12 or less

stillbirths per 1 000 total births resulting in an average global SBR of 9 deaths per 1 000 total births (other interim goals are shown in Fig. 5). All countries should focus on addressing inequalities and use audit data to track and prevent stillbirths.

Fig. 5 Ending preventable stillbirths



Source: The Lancet Every Newborn Series, Lawn JE et al. Lancet, 2014 (9).

The ambitious targets proposed in this action plan of ending preventable newborn deaths and stillbirths by 2035 – with intermediate targets for 2030, 2025 and 2020 – require universal, equitable and high-quality coverage

of essential, referral and emergency care for every woman and newborn in every country. This demands measurement, accountability and linkages with other global and national plans.





Strategic objectives and principles

To achieve the vision and goals, the Every Newborn action plan proposes five strategic objectives.



Strategic objective 1

Strengthen and invest in care during labour, birth and the first day and week of life.

A large proportion of maternal and newborn deaths and stillbirths occur within this period, but many deaths and complications can be prevented by ensuring high-quality essential care to every woman and baby during this critical time.



Strategic objective 2

Improve the quality of maternal and newborn care. Substantial gaps in the quality of care exist across the continuum for women's and children's health. Many women and newborns do not receive quality care even when they have contact with a health system before, during and after pregnancy and childbirth. Introducing high-quality care with high-impact, cost-effective interventions for mother and baby together – delivered, in most cases, by the same health providers with midwifery skills at the same time – is key to improvement.



Strategic objective 3

Reach every woman and newborn to reduce inequities. Having access to high-quality health care without suffering financial hardship is a human right. Robust evidence for approaches to ending preventable newborn deaths is available and, if applied, can effectively accelerate the coverage of essential interventions through innovations and in accordance with the principles of universal health coverage.



Strategic objective 4

Harness the power of parents, families and communities. Engaged community leaders and workers and women's groups are critical for better health outcomes for women and newborns. Education and empowerment of parents, families and communities to demand quality care and improve home care practices are crucial.



Strategic objective 5

Count every newborn through measurement, programme-tracking and

accountability. Measurement enables managers to improve performance and adapt actions as needed. Assessing outcomes and financial flows with standardized indicators improves accountability. There is a need to improve metrics globally and nationally, especially for birth outcomes and quality of care around the time of birth. Every newborn needs to be registered and newborn and maternal deaths and stillbirths need to be counted.

Principles

The Every Newborn action plan is based on six guiding principles.

Country leadership

Countries have primary ownership and responsibility for establishing good governance and providing effective and good-quality reproductive, maternal and newborn health services. Communities' participation in the planning, implementation and monitoring of policies and programmes that affect them is a central feature of such leadership and one of the most effective transformational mechanisms for action and accountability for newborn health. Development partners should align their contributions and harmonize actions.

Human rights

Principles and standards derived from international human rights treaties should guide all planning and programming for reproductive, maternal and newborn health and all phases of the programming process. Evidence and practice show the vital importance to health and development of many human rights outcomes.

Integration

Providing every woman and newborn with good-quality care that is available without discrimination and is accessible and acceptable requires integrated service delivery. Coordinated health system approaches involving multiple programmes, stakeholders and initiatives across the continuum of RMNCAH and nutrition are therefore essential, without losing visibility for newborn-specific content.

Equity

Equitable and universal coverage of highimpact interventions and a focus on reaching excluded, vulnerable and poorest population groups are central to realizing the rights of every woman and newborn to life, survival, health and development.

Accountability

Effective, accessible, inclusive and transparent programme-coverage and impact-monitoring mechanisms, independent review and action by all relevant actors are prerequisites for equitable coverage, quality of care and optimal use of resources. Accountability also includes access to processes and mechanisms for remedies, whether legal, administrative or other.

Innovation

Best practice evidence of strategies that broaden the coverage of interventions for newborns and reduce mortality has been accumulating over recent decades. Innovative thinking about ways to increase the participation of all stakeholders and reach the poorest and most underserved populations is nevertheless needed. More research and development is required to optimize the application of knowledge of which interventions and strategies are most effective.







Strategic objective 1:

Strengthen and invest in care during labour, birth and the first day and week of life

Rationale

The period occurring after 28 weeks of gestation to the first month after birth is especially important not just for survival, but also for early childhood interaction and development, when foundations for the evolution of cognitive and psychosocial skills are created. Furthermore, it is during labour, birth and the first week of life that 44% (1.2 million) of stillbirths, 73% (2 million) of newborn deaths and 61% (176 290) of maternal deaths occur (9).

Every pregnant woman should receive essential care provided by a skilled attendant who is proficient to monitor labour and assist the birth, able promptly to detect and manage complications competently, and capable of arranging for immediate referral when needed. Every baby should receive essential newborn care starting immediately after birth, during the first day, and continued at critical intervals in the first week of life and beyond.

Although globally the proportion of women giving birth with a skilled attendant (physician, nurse or midwife) has increased to 70%, great disparities in coverage and quality of care exist between and within countries. Coverage of skilled care at birth in sub-Saharan Africa reaches only half the population (29). Skilled

care around childbirth is most efficiently provided in many countries in public or private health facilities, as immediate access to emergency obstetric and newborn services when complications occur is crucial to the survival of mother and child.

Packages of proven interventions should ensure the provision of basic and additional care for women and newborns to prevent or treat the main causes of mortality. Providing extra care to small (either small for gestational age and/or preterm) and sick babies is particularly important in reducing neonatal mortality. Health personnel need to be sufficiently competent and equipped to support women and these babies, many of whom do not need advanced or intensive care and can be managed in a lower-level health facility or possibly in the community. Inpatient care facilities can play a vital role for babies who need full supportive facility care. Recent research indicates that simplified antibiotic regimens for treatment of possible serious bacterial infections delivered through outreach services from primary health facilities might save additional lives in settings where referral is not possible.

Research is an important element of investing in care around the time of birth and an integral



part of the actions proposed in this plan. Research priorities include understanding the factors that impede or facilitate extending coverage of proven interventions in lowand middle-income countries, ways to fill existing gaps (such as the need for greater understanding of the biological basis of term and preterm labour and new ways of preventing preterm birth) and investigation of the long-term effects, later in life, of morbidities occurring before and around conception, during pregnancy and in the first month of life. Improving data collection and fostering innovation and collaboration are also essential.

Proposed actions

Governments and all concerned stakeholders should raise awareness and foster recognition in communities that the time around childbirth and the first week of life is vital to saving maternal and newborn lives and assume accountability for creating the conditions in which every woman and newborn can realize their right to health and health care.

All countries are encouraged to develop or re-focus national strategies and action plans in line with the principles, goals, targets and strategic objectives of the action plan. Each government should conduct a systematic situational analysis and agree a set of core interventions and packages that match the

local context, are relevant to the burden of neonatal morbidity and mortality, and fit within the continuum of care. Equitable access to high-quality care during labour, childbirth and the first week of the postnatal period should be emphasized in all relevant country action plans.

Comprehensive maternal and newborn health

packages must be a part of core entitlements under existing and emerging universal health systems in all countries. National authorities should institute measures to increase coverage of skilled care at birth in health facilities. They should support the implementation of guidelines and policies to improve management during labour and childbirth, including the use of the partograph, a simple tool for monitoring labour and anticipating complications, and increase the number of postnatal visits to women and their babies. Where necessary, more midwives, auxiliary staff and community health workers should be trained and retained.

National authorities, supported by stakeholders, should cost plans and allocate sufficient funding for women's and children's health, with due emphasis on care around the time of birth and the first week of life. Governments and all concerned partners should ensure that investments in maternal and newborn health are sustained beyond 2015 and increased where needed.



Key actions for strategic objective 1: care at birth

Governments, in collaboration with stakeholders, should:

- a. conduct a systematic situational analysis and agree a core set of interventions and packages for the local context
- b. institute measures to increase the coverage of skilled care at birth in health facilities
- raise community awareness of the importance of the period around birth and the first week of life for preventing maternal and newborn deaths and stillbirths
- d. increase accountability of all relevant stakeholders
- e. develop or sharpen national plans for newborn health within the continuum of RMNCAH and nutrition
- f. allocate adequate financial resources to implement the national plan



Strategic objective 2:

Improve the quality of maternal and newborn care

Rationale

Skilled care at birth is provided by midwives, nurses and physicians in many countries, but women who give birth with a skilled attendant may receive suboptimal or poor-quality, nonrespectful care. The partograph is still not commonly used, and infection-prevention and newborn care practices immediately after birth are often harmful and do not adhere to the principles of the Baby-friendly Hospital Initiative, which include keeping the baby warm, keeping mother and baby together, initiating breastfeeding early and promoting exclusive breastfeeding. Creation of appropriate areas for newborn care is often overlooked. Increasing the number of births in health facilities is therefore not sufficient to reduce neonatal mortality, particularly early neonatal deaths. Quality care for mothers and newborns should be assured, even in natural calamities, disasters or emergencies.

It is estimated that one in three pregnant women needs some intervention during birth and between 5% and 15% require a Caesarean section (34). Many women and babies in higher-mortality settings will not have access to emergency obstetric, or even basic neonatal, care. The incidence of birth complications, intrapartum-related death and neonatal encephalopathy rises with increasing neonatal

mortality, reflecting lack of quality obstetric and neonatal emergency care. Though high Caesarean section rates are not desirable, rates of less than 5% are usually a marker of unavailability of maternal and neonatal emergency and intensive care.

Quality and equity of care affect health outcomes in lower-mortality settings. As it is difficult to predict need for emergency obstetric services, every maternity service should be able to provide basic life-saving interventions for women and newborns and have uninterrupted access to transport for referral when serious complications arise.

Quality of care is particularly important for reducing risks of disabilities or impairments. For example, preterm babies are vulnerable to eye complications. Blindness from retinopathy of prematurity is preventable by improving quality of neonatal care, including safer use of oxygen, and by detecting retinopathy early. Preterm infants must be followed up and their eyes must be checked.

A seamless continuum between primary care and referral-level facilities saves lives. Community-based skilled birth attendants may be common in low- and intermediate-mortality settings where human resources and capacity for training exist, but about one third of births



Quality and equity of care affect health outcomes for mothers and their babies

globally (mainly in higher neonatal mortality settings) occur at home without care from a health professional. Effective community-based approaches require a functioning continuum of care and effective linkages to health facilities with comprehensive emergency obstetric care.

There is overwhelming evidence that the standard of education of physicians, nurses and midwives is low in many countries. Midwifery curricula in some do not meet global standards, with students not acquiring the competences necessary to provide good-quality services with confidence. Limitations in regulation and professional association capacity mean that midwifery personnel have little legal protection and lack an organized voice to represent their interests. These factors, combined with staff shortages, poorly equipped facilities and low remuneration, lead to poor motivation and low quality of care.

Health services need to deal with risk factors for poor neonatal outcomes, such as adolescent pregnancy, short birth intervals, malnutrition (underweight and obesity), chronic disease (such as diabetes), infectious diseases (like tuberculosis and HIV disease), substance abuse (tobacco and alcohol use, for instance), domestic violence and poor psychological health. Workplace policies are important in supporting women during pregnancy and in the postnatal period and should include regulations to protect pregnant and lactating women from physically demanding work (35). Behavioural and community interventions to reduce exposure to potentially harmful pollutants, such as from traditional cook-stoves and secondhand tobacco smoke, are also necessary. Prevention, screening and management of sexually transmitted infections (such as HIV and syphilis), malaria and noncommunicable diseases are often implemented through specific programmes but have to be well integrated with maternal and newborn health services.

Existing programme platforms can provide specific links to improved newborn survival and

health and will contribute to strengthening the quality of provided health services, including those for family planning, HIV infection and tuberculosis, syphilis, malaria, water and sanitation, nutrition, integrated management of childhood illness, home visits in the postnatal period and immunizations (27).

Adolescent-friendly health services offering sexual and reproductive health need to be available to young people, as data show that adolescents are not reached by health services and HIV infection is increasing among them. Prevention of early and unintended pregnancy, along with care for girls and young women during pregnancy, birth and in the postnatal period, are vital to supporting their own and their babies' mental and physical health.

Many health facilities, particularly those in remote areas, do not have life-saving commodities for women and newborns, as identified by the United Nations Commission on Life-Saving Commodities (5). Challenges include unregistered new formulations of medicines, major supply-chain bottlenecks, stock-outs, costs in settings where services are not free, and carers and health care providers being uneducated about commodities and their effectiveness. The private sector's considerable expertise in developing, manufacturing and distributing medicines, medical devices and technologies must be harnessed to increase availability, especially in resource-poor settings.

Few global indicators exist for monitoring the quality of maternal and newborn care, and mechanisms for monitoring and evaluation are non-existent in many countries. Maternal and perinatal death surveillance and response can be a powerful approach to improving care quality. Many countries have adopted legislation requiring notification of maternal deaths, which can provide an entry point for a confidential enquiry into causes leading to maternal death and planning of remedial action. A similar approach could be taken to perinatal mortality.

Proposed actions

Governments, in collaboration with professional associations, academia, training institutions and other stakeholders, should regularly update national policies and guidelines for interventions around the continuum of care for women's and children's health relative to global evidence-based guidelines and locally defined strategies. Additionally, they should operationalize effective quality improvement systems and adopt and enforce the implementation of norms and standards for respectful and high-quality maternal and newborn care.

Governments should adopt competency-based curricula for training of health care workers and put in place regulatory frameworks defining the scope and practice of midwifery and nursing, including specific skills of caring for small-for-gestational-age or sick newborns, lactation counselling and support training, and the minimum standards of educational requirement needed. Rectifying the shortage of specialists, such as neonatologists and breastfeeding counsellors, should also be considered, where appropriate.

Staffing levels for each facility providing maternal and newborn care need to be planned in such a way that services can be provided on a continuous basis, 24 hours a day, seven days a week. Team work is essential: teams in first- and second-level referral hospitals should be multidisciplinary and include specialized obstetric, paediatric and anaesthetic staff to manage maternal and neonatal complications.

Maternity facilities must have appropriate infrastructure and be adequately equipped to provide the care needed by mothers and babies. The norms of infection prevention and biosafety must be respected. Electricity, water, sanitation and hand-washing facilities, clean toilets, appropriate spaces for women to give birth with privacy and dedicated areas to manage sick newborns safely must be in place.

After birth, women and newborns in health facilities should receive all essential services

before discharge. Mother and baby should be routinely accommodated in the same room with provision made for mothers to provide kangaroo mother care comfortably. Expression and storage of breast-milk should be encouraged in health facilities that care for preterm, small-for-gestational-age and/ or sick babies, complemented by milk banks in selected referral care facilities. Secondary and tertiary care facilities should have suitably equipped and staffed neonatal units and nurseries and be linked to primary care facilities through a well-functioning referral system. The required postnatal care visits in the first week after birth are necessary for counselling on health-seeking behaviours and for detection of any complications.

Life-saving commodities (including essential technologies) for women's and children's health should be included in every national essential medicines list and an uninterrupted supply chain to all facilities, especially the most peripheral, should be ensured.

Monitoring and improvement of quality of care must be instituted in all public and private maternity care services through, for example, maternal and perinatal death surveillance and response, birth and death registration (including fetal deaths or stillbirths) and periodic surveys of health facilities' availability and readiness. Information technology can assist in realtime monitoring: mobile phones, for example, are valuable in increasing communication and sharing data among health providers and communities. Community and service links aiming to improve care quality require investment; audit meetings and accountability are also key elements of the process of improving quality of care.

Motivation of staff is an important determinant for the quality of care. Innovative approaches should also be applied as appropriate, such as coaching, mentoring, accreditation and continuous education to improve access and quality of care. National authorities may consider incentives such as financial payments, bonuses and public recognition. Performance-

based financing is being introduced in several countries, but further investigation of its costeffectiveness and associated risks are needed.

Raising public awareness and increasing community involvement can accelerate improvements in quality of care. Parliamentarians, who represent voters, legislate, scrutinize and approve budgets and oversee government actions, are therefore seminal in determining women's and children's well-being. Civil society and local leaders, including business leaders, can strengthen political will and help to increase public awareness and community ownership of problems and solutions. A free flow of data and information is needed to enable this, with results from annual health sector reviews being made publicly available in line with recommendations made by the Commission

on Information and Accountability for Women's and Children's Health (4).

Engaging the private sector through publicprivate partnerships can bring multiple benefits, including: high-level advocacy; technology transfer to low-income countries; lower costs and increased availability of quality-certified essential medicines and medical devices; improved quality of care and the provision of evidence-based services by private practitioners; improved stewardship and regulatory function of governments; transport provision for emergency cases; stronger employer-based health services and workplace policies and programmes that support pregnant women and new mothers; and development of innovative technologies with the potential to reduce newborn and maternal deaths and prevent stillbirths and disability.



Key actions for strategic objective 2: improve quality of care

Governments, in collaboration with stakeholders, should:

- a. update national policies, guidelines, norms and standards for maternal and newborn care
- b. operationalize effective quality improvement systems for respectful, high-quality maternal and newborn care
- adopt competence-based curricula for training of health care workers, and put in place regulatory frameworks for midwifery and other health care personnel
- d. ensure postnatal care visits in the first week; and provision of quality extra care for small and sick newborns
- e. consider and evaluate innovative approaches to motivate staff and to improve access and quality of care; further investigate cost-effectiveness and associated risks of performance-based financing, as has been introduced in some countries

- f. ensure that all facilities are adequately staffed with multidisciplinary teams, are able to manage maternal and neonatal complications at referral-facility level, and have basic optimum infrastructure
- g. ensure life-saving commodities for women's and children's health are included in every national essential medicines list
- h. develop strategies to engage private-sector providers in increasing their advocacy for action on newborn and maternal deaths and stillbirths, and develop innovative technologies to improve newborn and maternal health outcomes
- i. institute maternal and perinatal death surveillance and response, including notification of maternal and perinatal deaths (preferably within 24 hours)
- j. enhance public oversight of the quality of maternal and newborn care through raising public awareness and increasing community involvement



Strategic objective 3:

Reach every woman and newborn to reduce inequities

Rationale

Every woman and newborn has the right to good-quality health care in line with the principles of universal health coverage and human rights. Access to high-quality maternal and newborn care depends, among other factors, on the availability of skilled health workers who are motivated, adequately equipped and equitably distributed. In many countries, inequitable access to health professionals, particularly in rural areas, is one of the main factors behind persistent high mortality rates for women and newborns.

Currently, fewer than one in six countries with the highest burden of maternal and neonatal mortality reaches the minimum benchmark of 23 doctors, midwives and nurses per 10 000 population necessary to provide a basic package of care (36). Severe shortages of midwives exist in at least 38 countries (37). These factors, combined with poor working conditions and few incentives for staff to live and work in remote areas or among disadvantaged populations, lead to unequal distribution of health workers and great inequities in access to care for mothers and newborns in countries with a high and inequitable burden of newborn deaths.

Costs of health services can present an important barrier to families seeking care during pregnancy and childbirth and in the postnatal period. Direct costs, such as over-

the-counter payments for medicines and fees for consultations and procedures, and indirect costs on, for example, transport and lost income have led to sharp inequities in coverage, most notably for women who give birth with a skilled attendant. Up to 11% of the population in some countries incur high costs in paying for health care, with as many as 5% forced into poverty because of health carerelated expenditure, including costs associated with essential maternal and newborn care (38). The goal of universal health coverage stipulates that everybody should be able to access health services and not be subject to financial hardship in doing so, but the world is falling short on both counts, particularly for women and children.

Special measures are necessary in health service and community settings to overcome violations of human rights where there is gender bias against the girl child.

The absence of information on budgets limits transparency and oversight for maternal and newborn health. Few countries have conducted national health accounts with specific subaccounts for maternal and newborn health, and tracking of development assistance has become prominent in the public domain only recently.

Private sector enterprises should take special measures to promote community and workplace support for mothers in relation to



Reaching all women and newborns requires investment in every aspect of the health system

pregnancy and breastfeeding and feasible and affordable child-care services, in compliance with the provisions of ILO's Maternity Protection Convention (35). The private sector can and should contribute to the promotion, protection and support of early and exclusive breastfeeding by ensuring that marketing and promotional practices fully conform to the provisions of the International Code of Marketing of Breast-Milk Substitutes (39) and subsequent relevant World Health Assembly resolutions (40) and by complying with national legal and regulatory provisions aimed at giving effect to them.

Reaching all women and newborns requires investment in every aspect of the health system, including leadership and governance, the workforce, infrastructure, commodities and supplies, service delivery, information systems and financing. Different contexts require tailored approaches, with specific attention to preparedness for, and rapid response to, complex humanitarian emergencies.

Proposed actions

A first proposed step for countries to move towards universal coverage of maternal and newborn care is to conduct a systematic analysis of the barriers to extending provision of the intervention packages necessary for provision of good-quality care. These can be found within the health system itself, but gaps in family and community knowledge, misperceptions and lack of resources can also contribute. Results of such analyses should be taken into consideration in the design of national newborn action plans.

Integration of actions on the health of newborns into existing RMNCAH initiatives and service delivery platforms (including those for HIV, malaria and immunization) will be crucial to ensuring that no opportunities to reach mothers and newborns are missed. Countries must critically assess what services are needed, motivate health workers, improve hospitals' efficiency, eliminate waste and corruption, and optimize the use of technologies and health services.

Every country should have a comprehensive development plan for the health workforce that covers education and training, the distribution, motivation and retention of skilled personnel, and redresses the mismatch between available health care workers and positions through a system that tracks from pre-service training through to posting and retention. It is essential that health workers have respectful attitudes and behaviours towards women and newborns, and working conditions should allow health workers to operate in a safe and respectful environment. Plans should include actions to expand and/or more equitably redistribute the workforce and, where needed, implement immediate remedial measures, including appropriate remuneration. Provision of housing, electricity and salary supplements has been effective in motivating health workers to serve in remote and underserved areas.

Optimizing performance of available staff is a priority. Evidence that life-saving interventions can safely be delegated to mid-level health workers is strong, but only if they are appropriately trained. Midwifery personnel, for example, should be able to provide the full range of emergency obstetric care where referral possibilities are limited.

Governments should adopt and enforce laws and policies on equity of access and quality of maternal and newborn care in public and private sectors. Policies should include: ensuring women's and children's universal access to health care services without enduring financial hardship; notification of maternal and perinatal deaths; context-specific approaches to HIV infection and infant feeding; ratification of ILO's Maternity Protection Convention (35); and legislation to implement the International Code of Marketing of Breast-Milk Substitutes (39).

Plans for extending maternal and newborn health services should be based on objective information and evidence. Countries need to estimate requirements for the number of pregnancies and births within a given setting and build and equip the necessary

infrastructure. They need to define a standard package of maternal and newborn care for each level of health provision and determine the number of staff and constitution of teams, including midwives and midwifery personnel and, where needed, specialists such as obstetricians, neonatal nurses and paediatricians. Geographical mapping of access points has been successful in several countries in supporting decision-making on the number and location of maternity facilities to reach the greatest number of families and communities. Sound planning should inform increased investment in care quality around childbirth.

Targeted programmes in and out of school are needed to expand adolescents and young adults' access to, and use of, modern contraceptive methods. Measures taken to prevent early pregnancies should include legislation to prevent or lower the number of girls who marry under the age of 18 years and life-skills education (for boys and girls) to help to keep girls in school, reduce rates of coerced sex, prevent early and unintended pregnancy before 20 years and inculcate positive cultural norms and traditions. Young people should be able to access contraceptives

and comprehensive sexual education that is empowering and which is designed to enable them to make sexual and reproductive decisions freely and responsibly. Countries should provide the enabling environment for this to happen. Youth-friendly health services should be available to give pregnant adolescents the full support they need to be well prepared for birth and parenthood, regardless of marital status.

Policies are needed to eliminate disparities in health care access, including subsidizing the cost of care and focusing on the most vulnerable population groups to ensure that good-quality maternal and newborn health services are available at an affordable cost at the point of use (which would mean free to many people). Countries should reduce reliance on out-of-pocket payments by increasing forms of prepayment with pooling of funds to share financial risks across the population. Prepayment typically involves taxes and other government charges and/or insurance. To mobilize resources, governments must prioritize the health sector in domestic budgets, increase efficiency in collection of national revenue and adopt innovative ways of raising funds through, for instance, taxes on tobacco and alcohol.



Countries must track total health expenditure by financial source (and per capita) and total expenditure on RMNCAH by financing source (and per capita). All major development partners should report their assistance for maternal and newborn health against their commitments and make covenants that would enable national authorities to establish predictable budgets and reinforce mutual accountability.

Community strategies to improve demand for services, birth preparedness and essential newborn care practices, including participatory women's groups and home visits supported by community health workers, volunteers or lay facilitators, have a critical role to play in reducing inequalities in mortality and access to care. Evidence suggests that participatory women's groups have the largest impacts on mortality among the poorest.



Key actions for strategic objective 3: reduce inequalities

Governments, in collaboration with stakeholders, should:

- a. conduct a systematic analysis of obstacles to achieving full-scale, high rates of coverage of effective intervention packages for quality care within the health system and community and take action to remove those obstacles
- b. ensure the integration of actions for newborn health into existing RMNCAH initiatives and service delivery platforms so that no opportunity to reach mothers and newborns is missed
- c. prepare and implement a development plan for the health workforce to ensure competency and respectful behaviour, improve the density, and increase motivation and retention of relevant health worker cadres as needed
- d. optimize performance of available staff by considering delegation of tasks to mid-level health personnel with appropriate training and support and the role of community health workers in bridging gaps between families and health services

- adopt and enforce laws and policies on equity of access and quality of maternal and newborn care in public and private sectors
- f. plan maternal and newborn health services on the basis of an evidence-based agenda, and staffing required for its delivery
- g. ensure elimination of barriers (social and financial) that limit access to care by mothers and newborns, including girls in some populations who are especially vulnerable
- h. track national health expenditures, including those for maternal and newborn health, and mobilize additional domestic resources
- i. develop targeted programmes in and out of school to expand adolescents' and young adults' access to, and use of, modern contraceptive methods and give pregnant adolescents the full support they need
- j. consider community strategies to improve demand for services, birth preparedness and essential newborn care practices, including home visits by community health workers and participatory women's groups



Strategic objective 4:

Harness the power of parents, families and communities

Rationale

It is vitally important to overcome barriers to accessing skilled care at birth and to harness the power of parents, families and communities, engaging them to seek care throughout pregnancy, birth and the first days and weeks of their children's lives. This is particularly important in low-income countries, where almost half of mothers do not receive skilled care during childbirth, more than 70% of babies born outside facilities receive no postnatal care and most maternal and newborn deaths and stillbirths occur (41). Many newborns die at home without any care having been sought.

It is essential to empower women, parents, families and communities to seek health care services when needed and to ensure they can provide recommended care in the home by themselves. Health outcomes, both positive and negative, are determined by decisions made within the household, the families' ability to reach care when needed and the quality of the services received when they arrive. Unfortunately, many actors outside of health services are often not engaged in discussions and efforts to improve their health and to increase the coverage of essential interventions.

Programmes that seek to strengthen health services through mobilizing community members to adopt healthy practices, shifting social norms to increase social support and addressing barriers to access have demonstrated the effects of such approaches.

The resulting changes are particularly evident for maternal and newborn health (42). A woman's right to make decisions is crucial for her health and the health of her family.

Community-oriented activities can be broadly categorized in four areas: (i) increasing awareness of rights, needs, responsibilities and potential problems related to maternal and newborn health; (ii) developing capacities to stay healthy, make healthy decisions and respond to obstetric and neonatal emergencies; (iii) strengthening social support networks among women, families and communities and links with the health services; and (iv) improving quality of care through strengthening health services' interactions with women, families and communities and responses to their needs (43). Investment is necessary in each of these areas, particularly in settings where maternal and neonatal mortality rates are high and access to health services is limited. Actions need to be taken collectively by multiple parties alongside efforts to improve the quality of health services. Civil society can play a catalytic role through existing and strengthened coalitions and networks.

Families, especially parents, are at the forefront of providing newborn care. They can and should ensure certain aspects of care for the healthy baby after birth, including keeping the baby clean and warm, initiating early and exclusive breastfeeding and performing proper cord care (including the use of chlorhexidine, if



Strengthening linkages between communities and health facilities improves health outcomes for women and their babies

appropriate). This is particularly important for babies born at home or discharged a few hours after birth. Even though practices like breastfeeding are often considered as natural behaviours, many women require skilled support.

Interactions between carer and child are crucial for psychosocial and cognitive development and should start from the day of birth. Simple communication and stimulation catalyse the maturation of neurological pathways and are relevant for term and preterm babies. Support for early child development is an essential component of newborn care.

Men can play an important role in maternal and newborn health as partners, fathers and community members. Men are often key decision-makers in maternal and newborn care-seeking behaviour and need to understand the needs, risks and danger signs of pregnancy, childbirth and postnatal periods. Health programmes that are traditionally designed to interact with women need to broaden their understanding of men's needs and perspectives (without compromising women's rights), working with men and women (adolescents and adults) and positioning gender perspectives and reproductive rights as pertinent to both. Health care workers need to make it convenient for men to accompany their partners and attend births, supporting them to enhance couples' communication and decision-making.

In 2009, WHO and UNICEF issued a joint statement on home visits for newborn children based on research showing that visits conducted by community health workers improved newborn survival rates (44). Extensive studies in Africa and Asia since then have shown that home visits during pregnancy and in the first week after birth increase the number of women seeking antenatal care and receiving skilled care during birth. Visits by community health workers also help families to take better care of the newborn and lead to improved practices, such as delayed bathing and early initiation of exclusive breastfeeding, resulting in significant reductions in neonatal mortality.

Community leaders, family members, traditional birth attendants and other influential people can also be positive agents for support (43). Depending on the context, elder family members, such as grandmothers and mothers-in-law, have a strong, culturally sanctioned power for decision-making and care, influencing, for example, the diet of pregnant women, workload and household responsibilities and use of emergency services.

Community mobilization through facilitated participatory learning and action with women's groups is recommended to improve maternal and newborn health, particularly in rural settings with low access to health services (45). The intervention can have a positive effect on newborn mortality, and further research may improve understanding of the effects on maternal health and care-seeking. Implementation of facilitated participatory learning and action with women's groups should focus on enabling discussions in which women can identify priority problems and advocate for local solutions for maternal and newborn health. In order to ensure quality, this intervention should be implemented with close monitoring and evaluation as well as prior adaptation to the local context.

Lay health workers, including traditional birth attendants, have successfully performed functions related to health care delivery for women and newborns and can be important members of the health team. They are not intended to replace a health professional in attending births, but can effectively promote maternal, newborn and reproductive health interventions such as, but not limited to. appropriate care-seeking, preparedness for birth and complications, and support for breastfeeding. Traditional birth attendants are often valued and respected community resources and finding new roles for them in areas such as providing continuous support for women during labour (in the presence of a skilled attendant) and serving as a link between communities and health services is important (46).

In conjunction with other actions, social and mass media can be influential in imparting knowledge, changing behaviour and instilling social accountability for newborn health services. Advocacy campaigns using radio and television have contributed to increasing the number of births in health facilities, early initiation and exclusive breastfeeding, and other interventions. Multi-pronged approaches enhance the reach of messages.

The private sector's considerable expertise in strategic communications can be harnessed to change social norms (such as acceptance that maternal and newborn deaths are inevitable), promote optimal health behaviours and increase demand for good-quality care. Many private corporations have capacity for conducting research and mining diverse sources of data to create understanding of what motivates behaviours, an extremely valuable asset that has not yet been tapped to its full potential. Public-private partnerships are especially amenable to multimedia advocacy campaigns through existing private sector communications platforms, television, radio, social media and e-health and mobile-telephone technologies.

Civil society organizations can contribute significantly to social mobilization, creating political will and policy design, and can help to hold governments and health services accountable for implementing their programmes and plans.

Systematic reviews of the results of evidence-based approaches designed to harness the power of women, parents and communities are being conducted in areas such as: preparedness for birth and complications; transport schemes; maternity waiting homes; community participation in programme planning and quality improvement; alleviation of financial barriers and cash transfers; integrated case management of newborn and childhood illnesses; the use of mobile telephone technology; maternal and perinatal deaths audits; and results-based financing (6). Findings will inform future policy on these approaches.

Proposed actions

Maternal and newborn deaths and stillbirths need to be recognized by communities, families and individuals as avoidable and not inevitable. To ensure zero tolerance of preventable deaths, there has to be a change in social norms and expectations surrounding childbirth and newborn survival, and parents affected by stillbirths or newborn deaths need to be given a voice. Peer-to-peer strategies of using parents' voices to mobilize civil society, mass media and social media can be used to spread information and change norms.

The analysis of obstacles proposed under strategic objective 3 should also focus on factors that determine the demand for maternal and newborn health services and affect the provision of recommended self-care in the home in countries with a high burden of neonatal mortality. Understanding the motivational, cultural, structural and financial elements that support families and communities to implement good home care practices and seek appropriate care for mothers and their newborns is essential for devising an appropriate response. Programmes should include activities to give women and families a voice and allow their participation in prioritizing problems and solutions.

Where access to health services is inequitable or low, countries should consider investing in community health workers as a powerful resource for improving maternal and newborn care, particularly in hard-to-reach areas, and ensure their integration as an important human resource alongside health professionals for maternal and child health matters. Appropriate training, supportive supervision, deployment and compensation are necessary.

Consideration should be given to strengthening community-based organizations such as women's groups to foster community mobilization through dialogue-based approaches such as facilitated cycles of participatory learning and action. This is particularly relevant among rural populations for whom access to care is limited. Further evaluation of the utility of this approach in poor

Community
mobilization
through
facilitated
participatory
learning and
action with
women's
groups can
improve
maternal
and newborn
health,
particularly in
rural settings

urban areas and with other actors, including men and grandmothers, is urgently needed.

Families, particularly men and fathers, also have important roles. They should be involved in individual household-level and community activities to enhance their contribution to supporting family health. Various channels can build capacity to provide good home care, such as health education, media campaigns and home visits by trained community health workers.

Strengthening linkages between communities and health facilities improves health outcomes for women and their babies, especially where referral services are provided. Mobile phone technology, which is now widely available and can reach remote areas in many countries, provides one approach for improving these linkages. It has been successfully used to supply health messages, establish help lines and facilitate real-time monitoring of births and deaths and can be linked with community mobilization interventions to create greater synergy.

Creating demand for services for underserved communities requires innovative approaches based on evidenced-based strategies. For example, conditional cash transfers to families and communities can be considered where financial circumstances impede access to health care, with careful monitoring to demonstrate cost–effectiveness.

The Commission on Information and Accountability for Women's and Children's Health has called for improved oversight and transparency, urging parliamentarians, community leaders, civil society and the general population to demand information and participate actively in planning and monitoring health services and the quality of care received by mothers and children. Countries should pursue these recommendations. Similarly, all concerned stakeholders should accept independent accountability for implementing this action plan at global level as part of their commitment to saving the lives of newborns.



Key actions for strategic objective 4: parents, families and communities

Governments, in collaboration with stakeholders, should:

- a. promote zero tolerance for preventable stillbirths and maternal and newborn deaths
- b. maximize the power of parents' voices, civil society, mass media and social media to provide information and change norms
- c. conduct a systematic analysis of obstacles
 to accessing quality maternal and
 newborn services and uptake of essential
 home care practices by women, families
 and communities and actively involve
 communities in determining priorities and
 planning appropriate actions
- d. equip families, including men, with the knowledge and capacities to provide good home care
- e. strengthen links between community and health facilities through applying innovative approaches to reach remote areas

- f. adopt combined approaches to improving care quality within the home and from health services
- g. evaluate the utility of community-based organizations such as participatory women's groups to foster community mobilization, particularly among rural populations with limited access to care
- adopt evidence-based strategies to generate and sustain demand for services using community-oriented actions
- i. engage, enable and support in-country civil society organizations to demand transparency and oversight and improve access to, and quality of, care
- j. engage the private sector to support multimedia communication campaigns to change social norms, promote zero tolerance for preventable mortality and advocate for optimal behaviours



Strategic objective 5:

Count every newborn – measurement, programme tracking and accountability

Rationale

Vital statistics provide indispensable information, in this case making policies more effective and responsive to the needs of women and children. In 2012, however, about one third of 137 million births globally and nearly all neonatal deaths and stillbirths went unregistered (9). Half the countries in the WHO African and South-East Asia regions do not record cause of death in their vital statistics, and serious deficiencies are present within existing systems (47). The vital registration systems in some countries do not follow global recommendations about which child to count and often function for only part of the country. In others, not all deaths are registered. Failure to collect high-quality data on registration of births and deaths, including cause of death, results in an absence of crucial information for policy-making, planning and evaluation across all development sectors, including health and health services. The United Nations Human Rights Council's resolution (48), on birth registration and the right of everyone to recognition everywhere as a person before the law, is entirely dedicated to birth registration and legal identification for all, without discrimination.

As governments and partners establish and expand access to interventions related to newborn health, more and better information is needed for monitoring and assessing progress towards achieving the commitments made to ending preventable newborn deaths and stillbirths.

Few universal indicators are available for monitoring equity of access and quality of maternal and newborn care. Some steps to improve measurement have been initiated, such as adding questions to household survey instruments on postnatal care for mothers and newborns and to facilityassessment tools about the availability of commodities specifically for newborns, including resuscitation equipment and antenatal corticosteroids. Many challenges remain, however, in gathering these data. Population-based household surveys, including demographic and health surveys and multiple indicator cluster surveys, have long collected data on indicators on family planning, antenatal care and attendance at birth. More information on postnatal contacts for newborns has recently been incorporated into these surveys and the number of countries with available data is steadily increasing. A process for generating



indicators that can be considered for addition to household surveys to measure newborn care practices and the content of postnatal care is a milestone proposed in this plan.

Major gaps persist in the collection of data on outcomes, coverage and quality of care around the time of birth, and much more rigorous attention needs to be paid to the development and testing of indicators and their inclusion in health management information systems. Few indicators related to the health of newborns are currently included in routine health management information systems, with limited use of the data for improving quality. A set of core and additional indicators for tracking not only population-based coverage of effective interventions, but also the quality of care in health services, needs to be agreed for use in varying contexts, including complex humanitarian emergencies. Managers at all levels need to know to what degree the system can deliver essential maternal and newborn services and identify performance weaknesses that can be rectified through better planning, budgeting and service delivery. Many of the indicators can be integrated in routine health information systems, with results validated periodically through specific surveys.

Many countries have accepted maternal death surveillance and response as an effective means of identifying deaths, investigating their determinants and taking remedial action on preventable causes. The introduction of perinatal audits in low- and middle-income countries has the potential to reduce mortality when solutions identified from the audit process were linked to action (49). New guidance from WHO and its partners provides clear recommendations on how to implement maternal death surveillance successfully at full scale (50). The guidance promotes a phased approach and suggests a focus on strengthening maternal death surveillance and response in health facilities before expanding it to communities. Auditing maternal and perinatal deaths and linking the results to a

national process has the potential to strengthen capacity to avoid preventable causes of mortality. Legal protection mechanisms that would facilitate full enquiries are nevertheless inadequate in many countries, meaning the full potential of the approach often remains untapped.

Proposed actions

Countries should introduce and invest in improving birth and death registration systems and consider innovative mechanisms for gathering data, such as through community health workers and use of mobile phones. Registration of stillbirths and newborn deaths should be accompanied by programmatically-relevant categorization of the causes of deaths. Additional data (such as gestational age and birth weight) and consideration of the social determinants of mortality should be included. The quality and completeness of data need to be monitored continuously and the data should be disseminated as the basis for planning.

Investment and technical support are needed to improve the quantity, consistency and frequency of national input data for all maternal and birth outcomes as part of a minimum perinatal dataset linked to vital registration and data derived from health facilities. Countries should strengthen mechanisms for maternal death surveillance and response and surveillance of perinatal mortality, linking this with perinatal death reviews and taking action to address avoidable factors identified through such reviews. It is also important to track disability outcomes (such as retinopathy of prematurity, deafness and cerebral palsy), particularly for countries expanding neonatal intensive care.

The global core indicators proposed as part of the action plan (Table 1) should be evaluated by stakeholders and, after assessment, integrated into national health information management systems. The set of indicators proposed will be further developed as part of the operationalization of the action plan, covering the domains of service readiness, outcomes, service use, intervention coverage and quality of care.

All stakeholders, from parliamentarians and community leaders to civil society and the general population, need to demand information and participate in planning and monitoring access to quality health services received by women and children, possibly through use of scorecards, to increase accountability for implementing this action plan.

While routine systems are being strengthened, countries and development partners should undertake periodic household surveys to obtain data on mortality, intervention coverage and use of services. The private sector should also be engaged in improving the collection and quality of birth and death registration systems and in investing, developing and executing innovative mechanisms for gathering data, especially through the use of mobile phones.

Key actions for strategic objective 5: count every newborn

Governments, in collaboration with stakeholders, should:

- a. invest in birth and death registration coverage and quality, promoting recording of every birth, live or stillbirth, and recording stillbirths and neonatal deaths
- b. consider the use of specific perinatal death certificates that capture additional data on stillbirths, gestational age and birth weight in addition to maternal complications into routine data collection systems and instruments

 g. urge parliamentarians, community leaders, civil society and the general population to
- c. develop a minimum perinatal dataset and ensure that all birth outcomes are collected, with consistent definitions and cross-links to databases for vital registration
- d. institutionalize maternal and perinatal death surveillance and response, linking this with perinatal death reviews and taking action to address avoidable factors identified through such reviews
- e. track morbidity and disability outcomes especially when neonatal intensive care is being expanded

- f. evaluate and define national indicators of service delivery for maternal and newborn health (based on the global indicators proposed in Table 1) and integrate them into routine data collection systems and instruments
- g. urge parliamentarians, community leaders, civil society and the general population to demand information and participate actively in planning and monitoring of access to quality health services received by women and children
- h. develop strategies to engage the private sector in improving the collection and quality of birth and death registration systems and in investing, developing and executing innovative mechanisms for gathering data




Framework for success

Impact framework

The impact framework shown in Fig. 6 inserts "Every Newborn" into the "Every Woman Every Child" concept and broadens the goals to include ending preventable deaths for women, stillbirths, newborns and children, and improving child development and human capital. The outcome level of the framework includes all packages for RMNCAH. This places the action plan's particular emphasis on care of women and babies during labour,

birth and the first week of life but also on care of small and sick newborns, with the objective of universal health coverage for all packages. Increased coverage of quality of care requires programme change to be expanded within the health system and community. Inputs to a strengthened health system require rigorous measurement, strong programme tracking and accountability. Finally, the socio-political, economic, environmental, biological and legal contexts affect all levels of change.

Fig. 6 Every Newborn impact framework



Mortality targets will only be achieved through improvements in coverage and quality of care for women and babies at birth and care of small and sick newborns. Interim targets of evidence-based interventions for coverage and

quality of care around birth, care for newborns at risk, home visits and participatory group support for women and newborns are therefore proposed (Fig. 7).

Every Newborn coverage targets by 2020

- a. Coverage and quality of care around birth: 90% of women giving birth and babies born in facilities will receive effective high-quality and respectful care that includes essential care during pregnancy, labour and following birth, with preventive care and appropriate management of complications for the mother and newborn. Maternal and perinatal death surveillance, timely response and regular monitoring of quality of care will be an integral part of maternal and newborn health services.
- b. Coverage and quality of care for small and sick newborns: at least half of babies who do not breathe spontaneously at birth after thorough drying and stimulation will be resuscitated with bag and mask ventilation; at least half of stable preterm newborns or babies weighing less than 2 000 g will receive kangaroo mother care and other supportive care; and at least half of newborns with possible serious bacterial infection will receive antibiotic therapy. Country-specific targets for comprehensive neonatal intensive care will be set, including tracking of disability.
- b. Home visits and participatory group support for women and newborns: each country will achieve at least a 20% increase (or an increase to 90% if their baseline is above 70%) of early postnatal care for women and newborns within two days of birth to promote breastfeeding, counselling and screening for maternal and newborn complications, and postnatal family planning. Linking to community participatory approaches and parent groups is an important component of this strategy.

Every Newborn coverage targets by 2025

- a. Coverage and quality of care around birth: 95% of women will give birth with a skilled attendant, and every woman and her newborn will receive effective high-quality and respectful care.
- b. Coverage and quality of care for newborns at risk: at least 75% of babies who do not breathe spontaneously at birth after thorough drying and stimulation will be resuscitated with bag and mask ventilation; at least 75% of stable preterm newborns or babies weighing less than 2 000 g will receive kangaroo mother care and other supportive care; and at least 75% of newborns with possible serious bacterial infection will receive antibiotic therapy.
- c. Home visits and participatory group support for women and newborns: 90% of women and newborns will receive early postnatal care of high quality within two days of birth. The quality of postnatal care will be tracked with improved metrics to assess content and longer-term outcomes, such as the nutrition goal of 50% exclusive breastfeeding in all countries by 2025. Linking to community participatory approaches and parent groups is an important component of this strategy.

Measures of success

The Every Newborn action plan is about taking action to achieve ambitious mortality targets and coverage targets to end preventable newborn deaths and stillbirths. The pathway to impact will be marked by milestones, which are defined at global and national levels for the period 2014–2020 (Fig. 7). The milestones will form the starting point for accountability and independent oversight and the basis for monitoring progress in implementation. Monitoring and evaluation coincide with the reviews of progress towards the MDGs in 2015 and will be ready for the prospective post-2015 sustainable development goals, linking to the new accountability mechanism.

A more detailed monitoring plan, with coverage and outcome metrics to track progress, is a milestone at global level. These indicators will need to be collected and used for national programme action within countries. Clearly delineating stillbirth interventions and strategies represents another milestone.

The real change for women and their babies will take place within countries. National milestones include (but are not limited to) ensuring that: commodities are included in national essential medicine lists and tracked; the community voice, especially of women, is heard; national health plans are sharpened and costed so as to deal appropriately with newborn health and stillbirths; and national expenditure for RMNCAH is tracked and reported.

Considering a woman and her baby together

forms a core concept of the action plan, so a package of interventions is needed for both: they are distinct yet interdependent, and the interdependence is vital to both. As a complement to this action plan, WHO, UNICEF and partners will develop an Every Mother Every Newborn Quality Initiative that will focus on improving quality of care for mothers and newborns in facilities, linked to community actions and district health system strengthening. A set of norms and standards for quality of care around birth and the immediate postnatal period will be defined and, once established in consultation with countries and technical experts, will need to be adapted and adopted within countries.

Achieving the vision, goals, and mortality and coverage targets outlined in the plan require measurable indicators to track progress and inform health policy and programmes. The selection of the core indicators (Table 1) involved a grading process for direct relevance to the action plan framework, targets and goals and review of current data availability. Core indicators in some cases are agreed and tracked, but some need to be further tested and integrated into national measurement systems. Urgent work is required to improve the metrics for these and other supporting indicators and increase the number of countries routinely tracking them. The operationalization of these core indicators and a wide list of additional necessary indicators forms part of the short-term milestones listed in Fig. 7.





GLOBAL LEVEL

2035

END PREVENTABLE NEWBORN DEATHS AND STILLBIRTHS

2035 Assessment of progress to national targets of 10 or less for newborn deaths and stillbirths

Global NMR target of 7 per 1000 livebirths and SBR of 8 per 1000 total births

2030

2030 Review of progress to national mortality targets of 12 or less for newborn deaths and stillbirths

Global NMR milestone of 9 per 1000 livebirths and SBR of 9 per 1000 total births

2025

2025 Review of progress to national mortality targets

Global NMR milestone of 12 per 1000 livebirths and SBR of 11 per 1000 total births

2020

2020 Review of progress to national mortality targets

Global NMR milestone of 15 per 1000 livebirths and SBR of 14 per 1000 total births

Global milestones by 2020

- Accountability in post-2015 plans: Ensure post-2015 development framework includes specific targets in newborn mortality and stillbirth reduction, in addition to under-5 child and maternal mortality reduction.
- Data: Monitoring plan, improving and using programmatic coverage data and equity, quality gap assessments, evaluation for improved indicators and investment to ensure that these are tracked at scale. Count every birth and death for women and babies including stillbirths, invest in civil registration and vital statistics, and innovate to improve and ensure the poorest are counted. Design and test a minimum perinatal dataset.
- Quality: Develop standards of quality and a core set of indicators for assessing quality of maternal and newborn care at all levels of health-care provision (Every Mother Every Newborn Quality Initiative).

- Investment: Ensure that investment in maternal and newborn health is continued in 2015 and sustained in the post-2015 development era.
- Innovation and research: Develop, adapt, and promote access to devices and commodities to improve care for mothers and newborn babies around the time of birth; and agree on, disseminate, and invest in a prioritzed and coordinated research agenda for improving preterm and newborn health outcomes.
 Particular focus is required for stillbirths, who have been left out and left behind.
- Coordination: Ensure coordinated support among UN partners, donors, academics, nongovernmental organizations and the private sector, and intensify efforts in the 20 countries that account for 80% of all newborn deaths.
- Champions: Develop new newborn champions, and engage champions for RMNCAH to integrate newborn messaging.

2014

NATIONAL LEVEL

2035 coverage targets assessment:

Universal coverage for all packages

2030 coverage targets assessment:

Achieve Sustainable Development Goals

- Care at birth:
 95% of births
 receive quality
 care
- 2025 coverage targets assessment:
- Care of small and sick newborn:
 >75% kangaroo mother care; >75% sepsis management; comprehensive neonatal intensive care: country-specific targets
- 3. Community care:90% coverage for postnatal care;50% exclusive breastfeeding at 6

2020 coverage targets assessment:

- Care at birth:
 90% of facility
 births receive
 high-quality care
- Care of small and sick newborns:
 >50% kangaroo mother care; >50% sepsis management; Comprehensive neonatal intensive care: country-specific targets
- 3. Community care:20% increase in postnatal care

National milestones by 2020

- National plans: Review and sharpen national strategies, policies, and guidelines for RMNCAH in line with the goals, targets, and indicators in *Every Newborn* action plan, including clear focus on care around the time of birth and small or sick newborn care.
- Data: Count every newborn by improving and using programmatic coverage data and equity, quality gap assessments. Institutionalise civil registration and vital statistics, adapt and use a minimum perinatal dataset, implement maternal/perinatal death surveillance and response.
- Quality: Adopt Every Mother Every Newborn Quality Initiative standards of quality and indicators for assessing quality of maternal/newborn care at all levels of health system; and ensure access to essential commodities for RMNCAH.
- Investment: Develop or integrate costed human resources for health strategy into RMNCAH plans, ensure sufficient financial resources are allocated.

- Health workers: Ensure the training, deployment, and support of health workers, in particular midwifery personnel, nurses, and community health workers.
- Innovation and research: Develop, adapt, and promote access to devices and commodities to improve care for mothers and newborn babies around the time of birth; and agree on, disseminate, and invest in a prioritized and coordinated research agenda for improving preterm and newborn health outcomes.
 Particular focus is needed for stillbirths, who have been left out and left behind.
- Engagement: Involve communities, civil society representatives, and other stakeholders to harness the power of individuals, families, and communities ensuring access and coverage of essential maternal and newborn care.
- Parent voices, champions: Shift social norms so that it is no longer acceptable for babies to die needlessly, just as it has become unacceptable for women to die giving birth.

Table 1 Proposed global indicators for the *Every Newborn* action plan

	Core Every Newborn indicators	Additional indicators
Impact	Maternal mortality ratio*	
	2. Stillbirth rate	Intrapartum stillbirth rate
	3. Neonatal mortality rate*	Low birth weight rate*
		Preterm birth rate
		Small for gestational age
		Neonatal morbidity rates, such as infection
		Rates of long-term disability after neonatal conditions
Coverage: care for all mothers and newborns	4. Skilled attendant at birth*	Early initiation of exclusive breastfeeding
	Exclusive breastfeeding for 6 months	_
	Early postnatal care for mother and babies*	
	7. Antenatal corticosteroid use	Caesarean section rate*
Coverage: complications and extra care	8. Newborn resuscitation	
	Kangaroo mother care and feeding support	
	10. Treatment of neonatal sepsis	
Input: counting	Birth registration*	Death registration, including cause of death
Input: Every newborn service delivery packages	Every Mother Every Newborn Quality Initiative with measurable norms and standards	
	Care of small and sick newborn	

 $^{^{*}}$ = indicator as defined and tracked in World Health Statistics 2013

Shaded = not currently routinely tracked

Bold = indicator requiring additional evaluation for consistent measurement and linked to milestones in Fig. 7

Italics = input package requiring norms and standards to be defined and linked to milestones in Fig. 7

All indicators are to be tracked in such a way that they can be broken down to assess equity by, for instance, urban/rural, regional, wealth quintile.



The way forward

Research priorities

Research into delivery, development and discovery needs to be placed at the forefront of efforts to reduce neonatal mortality. Research priorities for newborn health were identified by a global exercise for 2013–2025: nine out of 10 priorities related to improving delivery of known interventions.

The top research priorities for the delivery of interventions include: finding approaches to scale-up simplified newborn resuscitation at lower levels of the health system; identifying and managing newborn infection at community level; removing barriers to the extension of exclusive breastfeeding and facility-based kangaroo mother care; evaluating the use of chlorhexidine for cord care in neonates born in health facilities; and developing strategies to improve the quality of facility-based care provided during labour and childbirth.

Development research priorities identified included: adapting kangaroo mother care to make it deliverable at community level; detecting early high-risk women in pregnancy and labour in the community; improving and simplifying intrapartum monitoring; evaluating appropriate oral antibiotics for treatment of neonatal sepsis; defining the role of perinatal audits in improving quality of care during labour and childbirth; and developing lower-cost surfactant and devices for use in low- and middle-income countries.

Discovery research priorities highlighted: science and technology in order to understand the causal pathways of preterm labour; new tocolytics to delay preterm birth; stable surfactant with easier mode of delivery; effective maternal vaccines to prevent neonatal sepsis; point-of-care diagnostics and new biological agents better to identify and treat neonatal sepsis; better ways to detect fetal distress; and identification of biomarkers for intrauterine growth retardation and antepartum stillbirths.

Coordination

Putting the action plan into practice will require participation of many stakeholders, ranging from governments and policy-makers, donor countries, the United Nations and other multilateral organizations and global philanthropic institutions to civil society, health care workers and their professional associations, the business community, and academic and research institutions.

The plan defines priority actions to provide a healthy start for every newborn within the context of integrated RMNCAH programmes. Maternal and child health services are an ideal platform for delivering integrated packages that include a range of interventions, including those for malaria, HIV infection, nutrition and immunization. It is vital that post-2015 development goals include the vision of healthy societies in which women and adolescent girls, newborns and children survive and thrive. Coordination for implementation of this vision will rely on a strong, secure continuum of care to reduce dramatically preventable maternal, newborn and child deaths, and which has the potential to build more equitable societies and transform human development.

Actions by constituency

Governments and policy-makers at national, regional and global levels

- **National plans:** review and revise national strategies, policies and guidelines for RMNCAH in line with the goals, targets, principles and indicators defined in the *Every Newborn* action plan, including a clear focus on care around the time of birth.
- Budgets: allocate sufficient financial resources to maternal and newborn health, and ensure adequate investment to improve quality and equitable coverage of care.
- Legislation: adopt appropriate legislation on birth registration, maternal deaths notification, maternity protection and the International Code of Marketing of Breast-Milk Substitutes.
- Health workers: develop or integrate a costed strategy on human resources for health into RMNCAH plans to ensure the training, deployment and support of health workers, particularly midwifery personnel, skilled birth attendants, nurses and community health workers.

- Quality: adopt standards of quality and core set of indicators for assessing the quality of maternal and newborn care at all levels of health care provision.
- Commodities: include essential commodities for maternal and newborn health in national essential medicines lists and ensure an uninterrupted supply at all levels of the health system.
- Engage: engage with communities, civil society representatives, professional associations, the private sector and other stakeholders to harness the power of individuals, families and communities to ensure access and quality coverage of essential maternal and newborn care.
- Accountability: count every newborn by institutionalizing civil registration and vital statistics maternal, perinatal and neonatal death surveillance and response.

Organizations in the United Nations system and other multilateral agencies

- Policy: ensure that post-2015 development framework includes specific targets in newborn mortality reduction and stillbirth reduction, in addition to under-5 year old child mortality and maternal mortality reduction.
- Technical assistance: provide technical assistance and support to government planning, implementation and accountability efforts.
- Coordination: ensure coordinated support among United Nations partners and intensify efforts in the 20 countries that account for 80% of all newborn deaths.
- Quality: develop standards of quality and a core set of indicators for assessing quality of maternal and newborn care at all levels of health care provision.
- Investment: ensure that multilateral investment in maternal and newborn health is continued in 2015 and sustained in the post-2015 development era.
- Champions: engage champions for RMNCAH in order to provide coherent, coordinated and evidencebased messages about newborn health.

Donors and foundations

- Funding: mobilize funds to fill gaps and support the implementation of costed, evidence-based, countryowned RMNCAH plans that include a focus on birth.
- Health worker training: support the training and deployment of health workers, including investing in midwifery personnel, skilled birth attendants, nurses and community health workers that can deliver quality essential interventions focused on birth.
- Commodities: support access to quality commodities by investing in innovative financing, creating incentives for producers and purchasers, supporting quality assurance and regulation, and research and development efforts to improve products.
- **Accountability:** engage in country compacts and enhance accountability around financial flows.

Private business

- *Innovation:* invest in developing and adapting devices and commodities to care for mothers and newborns around the time of birth; invest in social and behavioural change campaigns, including those that reach the poorest and most vulnerable.
- Implement: scale-up best practices and partner with the public sector to improve and expand health worker training and quality service delivery.



Nongovernmental organizations, communities and/or parent groups

- Community health workers: support preventive care before and after the period around birth and referrals to basic and comprehensive facilities as appropriate.
- Community leadership and accountability: foster community leadership and accountability to remove barriers (in relation to, for instance, transport), hold health providers accountable for providing quality services and strengthen links between communities and facilities.
- Champions: identify and support local champions, including parliamentarians, parent groups, professionals, community health workers and community leaders; engage and link champions for RMNCAH in order to integrate coordinated and evidence-based messages about newborn health.
- **Academics and research institutions**
- Prioritize research needs: agree upon and disseminate a prioritized and coordinated research agenda for improving preterm and newborn health outcomes.
- *Invest in research:* encourage increased budget allocations for research into innovative interventions.
- **Health professionals**
- Essential interventions: prioritize essential interventions around the time of birth and care of small and sick newborns as part of an integrated package of RMNCAH services.
- Health workers: provide quality and respectful integrated services to babies and women through accelerated training, retention and motivation approaches.

- **Demand:** generate and sustain demand for services using community-owned actions (for instance, incentives such as conditional cash transfers, insurance, transport, social mobilization, savings credit schemes and cooperatives).
- Adolescents: give special attention to adolescent girls and implement approaches to help to prevent early and unintended pregnancies.
- Seek care: use families, communities and community health workers, skilled birth attendants and midwives in order to obtain essential maternal and newborn care that saves the lives of babies and women.
- Quality and accountability: be a voice for change; demand quality, affordable, accessible services; report poor services through government and nongovernment mechanisms.
- Build research capacity: build capacity at research institutions, especially in low- and middle-income countries, and train professionals.
- **Disseminate findings:** disseminate research findings and best practices.
- Build partnerships: strengthen global networks of academic providers, researchers and trainers.
- Commodities: work with local and national bodies to ensure consistent availability of commodities and supplies essential for key interventions around the period of birth.
- Quality: monitor quality of care, including through use of maternal and perinatal death surveillance and response.



References

- Darmstadt GL et al. Who has been caring for the newborn? Lancet, 2014, doi:10.1016/S0140-6736(14)60458-X.
- (2) Ban K. Global Strategy for Women's and Children's Health. New York: United Nations, 2010. (http://www.everywomaneverychild.org/images/content/files/global_strategy/full/20100914_gswch_en.pdf, accessed 12 June 2014).
- (3) A Promise Renewed. Committing to child survival: a promise renewed [website]. 2012 (http://www.apromiserenewed.org/A_ Promise_Renewed.html, accessed 24 May 2014).
- (4) Commission on Information and Accountability. Keeping Promises, Measuring Results: Commission on Information and Accountability for Women's and Children's Health. Geneva: World Health Organization, 2011.
- (5) Every Woman Every Child. UN Commission on Life-Saving Commodities for Women and Children [website]. United Nations Foundation; 2012 (http://everywomaneverychild. org/resources/un-commission-on-life-saving-commodities, accessed 27 March 2014).
- (6) Bhutta ZA et al. What will it take to avert preventable newborn deaths and stillbirths and at what cost? Lancet, 2014, doi:10.1016/S0140-6736(14)60792-3.
- (7) UN Inter-agency Group for Child Mortality Estimation (IGME). Levels and trends in child mortality: Report 2013. New York: UNICEF, 2013. (http://www.childinfo.org/files/Child_Mortality_ Report_2013.pdf, accessed 12 June 2014).
- (8) Dickson KE et al. Every Newborn: Health-systems bottlenecks and strategies to accelerate scale-up of newborn care in countries. Lancet, 2014, doi:10.1016/S0140-6736(14)60582-1.
- (9) Lawn JE et al. Every Newborn: Progress, priorities, and potential beyond survival. Lancet, 2014, doi:10.1016/S0140-6736/14/60496-7
- (10) Family Planning 2020 [website]. United Nations Foundation; (http://www.familyplanning2020.org/, accessed 2 February 2014).
- (11) Resolution WHA58.31. Working towards universal coverage of maternal, newborn and child health interventions. 2005. (http:// apps.who.int/iris/bitstream/10665/20381/1/WHA58_31-en. pdf, accessed 12 June 2014).

- (12) Stenberg K et al. Advancing social and economic development by investing in women's and children's health: a new Global Investment Framework. Lancet, 2013, 383(9925):1333-1354. doi: 10.1016/S0140-6736(13)62231-X.
- (13) Resolution WHA64.12. WHO's role in the follow-up to the United Nations High-level Plenary Meeting of the General Assembly on the Millennium Development Goals. 2010. (http://apps.who.int/gb/ebwha/pdf_files/WHA64-REC1/A64_REC1-en.pdf, accessed 12 June 2014).
- (14) Resolution WHA64.13. Working towards the reduction of perinatal and neonatal mortality. 2011. (http://apps.who.int/gb/ ebwha/pdf_files/WHA64/A64_R13-en.pdf, accessed 12 June 2014).
- (15) Resolution WHA64.12. Sustainable health financing structures and universal coverage 2011. (http://apps.who.int/gb/ebwha/ pdf_files/WHA64/A64_R9-en.pdf, accessed 12 June 2014).
- (16) United Nations. Convention on the Rights of the Child [website]. 1989 (http://www.ohchr.org/en/professionalinterest/ pages/crc.aspx, accessed 27 March 2014).
- (17) Office of the High Commissioner for Human Rights. Resolution A/HRC/21/L.10. Preventable maternal mortality and morbidity and human rights 2012. (http://www2.ohchr.org/english/ bodies/hrcouncil/docs/14session/A.HRC.14.39_AEV-2.pdf, accessed 12 June 2014).
- (18) WHO et al. Trends in maternal mortality: 1990-2013. Geneva: World Health Organization, 2014. (http://www.who.int/reproductivehealth/publications/monitoring/maternal-mortality-2013/en/, accessed 12 June 2014).
- (19) World Health Organization. Global Health Observatory Data Repository [website]. Geneva: World Health Organization; 2014 (http://apps.who.int/ghodata/, accessed 27 March 2014).
- (20) March of Dimes et al. Born Too Soon: The Global Action Report on Preterm Birth. Geneva: World Health Organization, 2012.
- (21) Lee AC et al. National and regional estimates of term and preterm babies born small for gestational age in 138 lowincome and middle-income countries in 2010. The Lancet Global Health, 2013, 1(1):e26-e36. doi:10.1016/S2214-109X(13)70006-8.

- (22) Wall SN et al. Neonatal resuscitation in low-resource settings: what, who, and how to overcome challenges to scale up? International Journal of Gynaecology and Obstetrics, 2009, 107 Suppl 1:S47-62, S63-44.
- (23) Blencowe H et al. Preterm-associated visual impairment and estimates of retinopathy of prematurity at regional and global levels for 2010. Pediatric research, 2013, 74 Suppl 1:35-49. 10.1038/pr.2013.205.
- (24) Resolution WHA63.17. Birth defects. 2010. (http://apps.who.int/gb/ebwha/pdf_files/WHA63/A63_R17-en.pdf, accessed 12 June 2014).
- (25) World Health Organization. Compilation of WHO recommendations on maternal, newborn, child and adolescent health [website]. Geneva: World Health Organization; 2014 (http://www.who.int/maternal_child_adolescent/documents/ mnca-recommendations/en/index.html, accessed 27 March 2014).
- (26) The Partnership for Maternal Newborn Child Health. A Global Review of the Key Interventions Related to Reproductive, Maternal, Newborn and Child Health (RMNCH). Geneva: PMNCH, 2011. (http://www.who.int/pmnch/topics/part_ publications/essential_interventions_18_01_2012.pdf, accessed 12 June 2014).
- (27) Mason EM et al. From evidence to action to deliver a healthy start for the next generation. Lancet, 2014, doi:10.1016/S0140-6736(14)60750-9.
- (28) World Health Organization. WHO recommendations:

 Postnatal care of mother and newborn. Geneva: World Health
 Organization, 2013. (http://www.healthynewbornnetwork.org/
 sites/default/files/resources/Postnatal_Care_Guidelines_web_
 v2.pdf, accessed 12 June 2014).
- (29) World Health Organization. World Health Statistics 2013. Geneva: World Health Organization, 2013. (http://www.who.int/gho/publications/world_health_statistics/2013/en/, accessed 12 June 2014).
- (30) World Health Organization. Preventing early pregnancy and poor reproductive outcomes among adolescents in developing countries. Geneva: World Health Organization, 2011. (http:// www.who.int/maternal_child_adolescent/documents/ preventing_early_pregnancy/en/, accessed 12 June 2014).
- (31) UNFPA. The State of the World Population 2013: Motherhood in Childhood: Facing the challenge of adolescent pregnancy. New York: United Nations Population Fund, 2013. (http://www.unfpa.org/webdav/site/global/shared/swp2013/EN-SWOP2013-final.pdf, accessed 12 June 2014).
- (32) Gogia S, Sachdev HS. Home visits by community health workers to prevent neonatal deaths in developing countries: a systematic review. Bulletin of the World Health Organization, 2010, 88(9):658-666B. doi: 10.2471/BLT.09.069369.
- (33) Kirkwood BR et al. Effect of the Newhints home-visits intervention on neonatal mortality rate and care practices in Ghana: a cluster randomised controlled trial. Lancet, 2013, 381(9884):2184-2192. doi: 10.1016/S0140-6736(13)60095-1.
- (34) WHO et al. Averting Maternal Death and Disability: monitoring emergency obstetric care. A handbook. Geneva: World Health Organization, 2009. (http://www.who.int/reproductivehealth/ publications/monitoring/9789241547734/en/, accessed 12 June 2014).
- (35) World Health Organization. Global Health Atlas of the Health Workforce. Geneva: World Health Organization. 2014. (http://apps.who.int/globalatlas/default.asp, accessed 12 June 2014).
- (36) UNFPA. State of the World's Midwifery Report 2011: Delivering health, saving lives. New York: UNFPA, 2011.

- (37) World Health Organization. World Health Report 2010: Health systems financing: the oath to universal coverage Geneva: World Health Organization, 2010. (http://www.who.int/ whr/2010/en/, accessed 12 June 2014).
- (38) International Labour Organization. C183 Maternity Protection Convention, 2000 (no. 183) [website]. 2000 (http://www.ilo.org/dyn/normlex/en/f?p=NORMLEXPUB:12100:0::NO::P12100_ILO_CODE:C183, accessed 27 March 2014).
- (39) World Health Organization. International Code of Marketing of Breast-Milk Substitutes. Geneva: World Health Organization, 1981. (http://www.who.int/nutrition/publications/ infantfeeding/9241541601/en/, accessed 12 June 2014).
- (40) Resolution WHA61.20. Infant and young child nutrition: biennial progress report. 2008. (http://www.who.int/nutrition/topics/ WHA61.20_iycn_en.pdf?ua=1, accessed 12 June 2014).
- (41) Requejo JH et al. Accountability for maternal, newborn and child survival: The 2013 Update. Geneva: World Health Organization and UNICEF, 2013.
- (42) Lee AC et al. Linking families and facilities for care at birth: what works to avert intrapartum-related deaths? International Journal of Gynaecology and Obstetrics, 2009, 107 Suppl 1:S65-85, S86-68.
- (43) WHO et al. WHO-UNICEF Joint Statement on home visits for the newborn child: a strategy to improve survival. Geneva: World Health Organization, 2009 April 11, 2014. (http:// www.healthinternetwork.com/maternal_child_adolescent/ documents/who_fch_cah_09_02/en/, accessed 12 June 2014).
- (44) World Health Organization. Working with individuals, families and communities. Geneva: World Health Organization, 2003. (http://www.who.int/reproductive-health/publications/ RHR_03_11/, accessed 12 June 2014).
- (45) World Health Organization. WHO Recommendation on community mobilization through facilitated participatory learning and action cycles with women's groups for maternal and newborn health. Geneva: World Health Organization, 2013
- (46) World Health Organization. Optimizing Health Worker Roles for Maternal and Newborn Health. Geneva: World Health Organization, 2012. (http://www.who.int/workforcealliance/media/news/2012/whotaskshiftingrecom/en/index.html, accessed 12 June 2014).
- (47) Oomman N et al. Modernising vital registration systems: why now? Lancet, 2013, 381(9875):1336-1337. doi:10.1016/ S0140-6736(13)60847-8.
- (48) Resolution A/HRC/19/L.24. Birth registration and the right of everyone to recognition everywhere as a person before the law. 2012. (http://www.ohchr.org/Documents/HRBodies/ HRCouncil/RegularSession/Session19/A-HRC-19-2_en.pdf, accessed 12 June 2014).
- (49) Pattinson R et al. Perinatal mortality audit: counting, accountability, and overcoming challenges in scaling up in low- and middle-income countries. International Journal of Gynaecology and Obstetrics, 2009, 107 Suppl 1:S113-121, S121-112.
- (50) World Health Organization. Maternal death surveillance and response: technical guidance. Information for action to prevent maternal death. Geneva: World Health Organization, 2013. (http://www.who.int/maternal_child_adolescent/documents/ maternal_death_surveillance/en/, accessed 12 June 2014).

Summary of action plan process

Although remarkable progress has been made in recent decades to reduce the number of child deaths worldwide, the NMR globally has declined at a slower pace despite a large proportion of newborn deaths being preventable. Opportunities for improving newborn health are unprecedented as, today, much more is known about effective interventions and service delivery channels, and approaches to accelerating coverage and raising quality of care. Recently, renewed commitments to saving the lives of newborn infants have been made by many governments and partners, in response to the United Nations Secretary-General's Global Strategy for Women's and Children's Health and its accompanying *Every Woman Every Child* initiative and to recommendations made by the Commission on Information and Accountability for Women's and Children's Health and the United Nations Commission on Life-Saving Commodities. This response has triggered an initiative by multiple stakeholders to propose to the global health community the development of a draft global action plan.

Preparation of the draft action plan

The preparation of the newborn health draft action plan has been guided by the advice of experts and partners, led by WHO and UNICEF, and by the outcome of several multi-stakeholder consultations at different global and regional forums.

A further consultative process followed with Member States, in particular regarding the goals and actions being proposed for the five strategic directions of the draft action plan and their related targets, through one global and two regional meetings of stakeholders. In addition a draft of the action plan was posted on the WHO website in December 2013 for consultation by Member States and other stakeholders before the 134th session of the Executive Board. At that session, the Board noted the proposed process for the further development of the draft action plan.

The subsequent web-based consultation was conducted, as outlined in document EB134/17 Add.1. A Note Verbale was sent to Member States on 4 February 2014, describing the process and inviting feedback. In addition, information about the web consultation was widely disseminated through social media and networks on RMNCAH to encourage as broad a response as possible.

By the deadline of 28 February 2014, more than 300 comments had been received, including responses from 43 State, actors, 23 professional associations, 102 nongovernmental organizations, and many individuals. The Secretariat took these comments into account in revising the draft action plan. The final plan was endorsed at the 67th World Health Assembly in May 2014 and supported by resolution 67/14.2 (Annex 1). Member States made firm commitments to its implementation and requested the Director General to monitor progress towards achievement of the global goal and targets and report periodically to the Health Assembly until 2030.

The proposed "Every Newborn: an action plan to end preventable deaths", provides clear objectives and actions for Member States and other stakeholders. It also proposes indicators that can be used to evaluate both progress in implementation and the impact of the action plan. A detailed monitoring plan with coverage and outcome metrics to track progress towards the goal and targets will be published as a separate document.





Annex 1

Sixty-Seventh World Health Assembly Resolution WHA67.10: Newborn health action plan



SIXTY-SEVENTH WORLD HEALTH ASSEMBLY Agenda item 14.2

WHA67.10 24 May 2014

Newborn health action plan

Having considered the reports on the newborn health: draft action plan,¹ monitoring the achievement of the health-related Millennium Development Goals,² and health in the post-2015 development agenda;³

Recalling resolution WHA58.31 on working towards universal coverage of maternal, newborn and child health intervention, resolution WHA63.15 on monitoring of the achievement of the health-related Millennium Development Goals, resolution WHA64.9 on sustainable health financing structures and universal coverage, resolution WHA64.13 on working towards the reduction of perinatal and neonatal mortality, and resolution WHA65.7 on implementation of the recommendations of the Commission on Information and Accountability for Women's and Children's Health;

Acknowledging the pledges and commitments made by a large number of Member States and partners to the United Nations Secretary-General's Global Strategy for Women's and Children's Health, which aims to save 16 million lives by 2015;

Recognizing that millions of children and women die needlessly each year during and around the time of childbirth, and that effective interventions are available and feasible for implementation at scale to end preventable maternal, newborn and child deaths;

Recognizing that ending preventable maternal mortality will accelerate the achievement of the newborn mortality target;

Concerned that there has been insufficient and uneven progress towards achieving Millennium Development Goal 5 (Improve maternal health);

Also concerned that, although progress has been made towards achieving Millennium Development Goal 4 (Reduce child mortality) in terms of the overall reduction of child mortality, the reduction of perinatal and neonatal mortality has stagnated and the proportion of neonatal deaths among all child deaths is increasing;

Recognizing the need to intensify action urgently in order to end preventable neonatal deaths and preventable stillbirths, especially by improving access to and quality of health care for women and newborns, particularly of those at risk, especially for high-risk groups and including the prevention of the transmission of HIV from mother to child, within the continuum of care for reproductive, maternal, newborn and child health,

¹ Document A67/21.

² Document A67/19.

³ Document A67/20.

WHA67.10

- 1. ENDORSES the newborn health action plan,1
- 2. URGES Member States² to put into practice the newborn health action plan, through steps
 - (1) reviewing, revising and strengthening their national strategies, policies, plans and guidelines for reproductive, maternal, newborn and child health in line with the goal, targets and indicators defined in the newborn health: action plan, and strongly committing to their implementation with particular focus on high-risk groups;
 - (2) committing themselves, according to their capacities, to allocating adequate human and financial resources to improve the access to and the quality of care, particularly care for the mother and the newborn during labour, around birth and the first week, and achieve the national newborn health targets in line with the global action plan;
 - (3) strengthening health information systems so as better to monitor quality of care and to track progress towards ending preventable maternal and neonatal deaths and stillbirths;
 - (4) sharing information on lessons learnt, progress made, remaining challenges and updated actions to reach the national newborn and maternal health targets;
- 3. REQUESTS the Director-General:
 - (1) to foster alignment and coordination of all stakeholders to support the implementation of the newborn health action plan;
 - (2) to identify and mobilize, within approved current and subsequent programme budgets, more human and financial resources for the provision of technical support to Member States in implementing the newborn health component of national plans and monitoring their impact;
 - (3) to prioritize the finalization of the more detailed monitoring plan with coverage and outcome metrics to track progress of the newborn health action plan;
 - (4) to take into due account the views expressed at the Sixty-seventh World Health Assembly as well as the domestic context when supporting the implementation of the action plan at the national level:
 - (5) to monitor progress and report, periodically until 2030, to the Health Assembly on progress towards achievement of the global goal and targets using the proposed monitoring framework to guide discussion and future actions.

Ninth plenary meeting, 24 May 2014 A67/VR/9

= = =

¹ "Every newborn: an action plan to end preventable deaths" contained in document A67/21.

² And, where applicable, regional economic integration organizations.

Annex 2

Ending Preventable Maternal Mortality (EPMM) – Targets beyond 2015

A series of technical consultations and discussions convened by WHO, UNFPA, UNICEF, USAID, the Maternal Health Task Force (MHTF), and the Maternal and Child Health Integrated Program (MCHIP) and attended by a large number of stakeholders has resulted in the proposal of a vision for ending preventable maternal mortality (EPMM) and maternal mortality targets for 2030. The latest of these consultations, held in April 2014 in Bangkok, was attended by over 95 participants from 34 countries, including many countries with high rates of maternal mortality. At this consultation it was affirmed that EPMM is within reach and that progress can be accelerated by positioning maternal survival in the context of every woman's right to healthcare and the highest attainable level of health across the lifespan. In acknowledgement that maternal health is a crucial element of development, the consultation forged consensus on maternal mortality reduction targets for inclusion in the post-2015 development agenda.

Targets for equitable GLOBAL maternal mortality reduction post-2015

Global target: Reduce global Maternal Mortality Ratio (MMR) to less than 70 maternal deaths per 100,000 live births by 2030.

Secondary global target: By 2030, no country should have an MMR greater than 140, a number twice the global target.

Achieving the average global target implies that all countries contribute by reducing their MMR in 2010 by at least two-thirds by 2030. In order to meet the global target, countries with the highest MMRs (MMR >420) will need to reduce their MMR at an annual rate of reduction greater than 5.5%. Therefore, the secondary target aims to eliminate extremes of inequity in global maternal survival.

COUNTRY targets to reduce inequity in global MMR reduction

For countries with MMR less than 420 in 2010 (the majority of countries worldwide): Reduce the MMR by at least two-thirds from the 2010 baseline by 2030.

For all countries with baseline MMR greater than 420: The rate of decline should be greater, and in 2030, no country should have an MMR over 140.

To address inequity, the global community should work together to reduce extremes of mortality and avert a situation where some countries have MMRs far above the global average and eliminate disparities in MMR between sub-populations within countries.

Photo credits

Front cover photo: @Kiselev Andrey Valerevich/shutterstock.com

Page 4 photo: ©Isabel Pinto/PMNCH Page 5 photo: ©Isabel Pinto/PMNCH

Page 8 photo left: ©UNICEF/NYHQ1994-0632/Dora Gutierre
Page 8 photo right: ©UNICEF/NYHQ2005-1225/Roger LeMoyne

Page 9 photo: ©Jon Warren/World Vision

Page 11 photo: @Seifu Assegid/Save the Children

Page 14 photo: @UNICEF/NYHQ1993-0268/Roger LeMoyne

Page 18 photo: ©WHO/Jim Holmes

Page 19 photo left: ©UNICEF/PAKA2009-4180/Marta Ramoneda Page 19 photo right: ©UNICEF/AFGA2007-00097/Anita Khemka

Page 21 left photo: ©Ratana Lay/World Vision

Page 21 right photo: @UNICEF/MLWB2009-00040/Giacomo Pirozzi

Page 22 photo: ©lsabel Pinto/PMNCH

Page 24 photo: @Ayesha Vellani/Save the Children

Page 28 photo: @UNICEF/BANA2006-00503/Shehzad Noorani

Page 30 photo: @UNFPA/Edward Echwalu

Page 32 photo: @UNICEF/Pakistan/Laura Salvinelli

Page 36 photo: @UNICEF/NYHQ1993-0407/Roger LeMoyne

Page 41 left photo: @March of Dimes
Page 41 right photo: @Isabel Pinto/PMNCH

Page 45 photo: @March of Dimes

Page 47 photo: ©Shafiqual Alam Kiron/Save the Children

Page 48 photo: @March of Dimes

Page 50 photo left: @Sven Torfinn/Panos/H4+

Page 50 photo right: @UNICEF/UGDA2010-00754/Shehzad Noorani

Page 54 photo: @Jane Hahn/Save the Children



Acknowledgements

Every Newborn Steering Committee: Martina Lukong Baye, Ministry of Health, Cameroon; Anthony Calibo, Department of Health, Philippines; Luc de Bernis, United Nations Population Fund; Joseph de Graft Johnson, Maternal and Child Health Integrated program; Cyril Engmann, Bill and Melinda Gates Foundation; Suzanne Fournier, Children's Investment Fund Foundation; Lily Kak, USAID; Eve Lackritz, Global Alliance to Prevent Prematurity and Stillbirth; Gillian Mann, DFID; Charles Mwansambo, Ministry of Health, Malawi; Adetokunbo Oshin, Minsitry of Health, Nigeria; Vinod Paul, All India Institute of Medical Sciences; Yogan Pillay, Department of Health, South Africa; Joy Riggs-Perla, Save the Children, Saving Newborn Lives; Aline Simen Kapeu, UNICEF.

Writing Team: Mary Kinney (managing editor), Bernadette Daelmans, Kim Dickson (Chair), Joy Lawn, Elizabeth Mason (Chair), Lori McDougall, Juana Willumsen, Severin von Xylander.

In addition, we are grateful for the support and review of the individuals from the following countries, as members of the Advisory Group: Gloria Quansah Asare, Family Health Division, Ghana; Anuradha Gupta, Ministry of Health and Family Welfare, India; Maria Paulina Giusti, Ministry of Health, Peru; Ewlyn Chomba, Department of Health, Zambia.

More than 800 interested stakeholders participated in country bottleneck analysis workshops, country and regional meetings and reviewed the draft action plan, in addition to them, over 300 individuals, organizations and state actors who submitted comments to the online consultation process. A list of organizations that participated in this process is available at http://www.who.int/maternal_child_adolescent/topics/newborn/enap_consultation/en/.

We acknowledge and thank all these dedicated stakeholders for their valuable contributions to the development of the action plan.

Communications, advocacy, editorial and technical support: Kel Currah, Ian Hurley, Nathalie Likhite, Alex Mathieson, Veronic Verlyck, Linda Vesel.

Production coordinator: Gael Kernen.

Design and layout: GMMB, USA; Paprika, France.

Media: GMMB.

More information

Every Newborn **www.everynewborn.org**Healthy Newborn Network **www.healthynewbornnetwork.org**

Recommended citation

WHO, UNICEF. 2014. Every Newborn: an action plan to end preventable deaths. Geneva: World Health Organization.



























Advisory group































































