


How to close the maternal and neonatal sepsis gap in sub-Saharan Africa

Akaninyene Otu,¹ Emmanuel Fru Nsutebu,² Jane E Hirst,³ Kelly Thompson,⁴ Karen Walker,⁴ Sanni Yaya ^{5,6}

To cite: Otu A, Nsutebu EF, Hirst JE, *et al.* How to close the maternal and neonatal sepsis gap in sub-Saharan Africa. *BMJ Global Health* 2020;5:e002348. doi:10.1136/bmjgh-2020-002348

Handling editor Seye Abimbola

Received 28 January 2020

Revised 5 April 2020

Accepted 7 April 2020



© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Department of Infection and Travel Medicine, Leeds Teaching Hospitals NHS Trust, Leeds, United Kingdom

²Division of Infectious Diseases, Sheikh Shakhboub Medical City, Abu Dhabi, United Arab Emirates

³Nuffield Department of Women's and Reproductive Health, University of Oxford, Oxford, United Kingdom

⁴The George Institute for Global Health, University of New South Wales, Sydney, New South Wales, Australia

⁵School of International Development and Global Studies, University of Ottawa, Ottawa, Ontario, Canada

⁶The George Institute for Global Health, University of Oxford, Oxford, Oxfordshire, United Kingdom

Correspondence to

Dr Sanni Yaya;
sanni.yaya@uottawa.ca

INTRODUCTION

Sepsis is life-threatening organ dysfunction caused by a dysregulated host response to infection.¹ The incidence of sepsis, as estimated by the Global Burden of Disease study, was 48.9 million cases with 11 million deaths in 2017.² In sub-Saharan Africa (SSA), the burden is highest, with an estimated 16.7 million cases and sepsis accounting for up to 45% of all deaths, in many SSA countries.²

Maternal sepsis is a sepsis that occurs during pregnancy, childbirth, postabortion or in the postpartum period and neonatal sepsis is a sepsis that occurs within the first 28 days of life. Reliable estimates of the incidence and mortality of maternal and neonatal sepsis in SSA are incomplete. Notwithstanding this limitation, the most recent estimate suggests that sepsis is responsible for 130 000 maternal deaths³ and 300 000 neonatal deaths annually, though this may be an underestimate,⁴ in view of the recent Global Burden of Disease study report.²

Sepsis-related deaths in SSA reflect underlying political, poverty, health inequity and health system challenges across Africa.⁵ These challenges affect preventative and public health efforts to prevent sepsis and limit prompt and effective recognition and treatment of sepsis to prevent mortality. The challenges include access to clean water and sanitation, vaccination and provision of primary care in addition to access to antimicrobial agents and strategies for their stewardship.⁶⁻⁹

In 2017, the World Health Assembly (WHA) passed a resolution to improve the prevention, diagnosis and management of sepsis (WHA 70.7).¹⁰ This resolution recommended key action areas for member states and future priorities.¹¹ Based on these recommendations, we outline nine key areas to accelerate progress in reducing death due to maternal and neonatal sepsis in SSA.

HOW TO REDUCE MATERNAL AND NEONATAL SEPSIS IN SSA

1. Increase sepsis awareness in the community

Despite sepsis being a major cause of health loss globally, public awareness remains low,¹¹ with the delayed time from presentation to care in low-middle income countries (LMICs) contributing to adverse outcomes.^{12,13} Increasing awareness among the public and health workers is vital for prevention. In high-income countries (HICs), strategies to increase public awareness include public service announcements of the early warning signs of sepsis, published in traditional print and electronic media.¹⁴ With global mobile connections standing at 7.6 billion¹⁵ and extremely high smartphone dissemination in SSA, there is potential for exploitation of this digital resource to increase sepsis awareness in LMICs. Rapid evaluation of public awareness can be measured using geospatial analysis of online information-seeking behaviours as a surrogate measure.¹⁶

Educational messages can be accessed remotely and in real-time on mobile phones by health workers in different geographic locations, unrestricted by time and space. Mobile applications (apps) are conceptualised as a set of tools designed to undertake specific tasks and jobs. A systematic review of 42 studies to assess the effectiveness of mobile health in LMICs supports their use.¹⁷ The use of mobile phone apps in supporting, training, data collection and reporting, emergency referrals, work planning and communication between health workers was associated with improvements in adherence to treatment algorithms, more accurate data collection and a reduction in treatment errors.¹⁷

2. Improve sepsis surveillance networks

Sepsis surveillance systems are critical tools for comprehensively quantifying the burden of sepsis related morbidity and mortality over time. At facility level, surveillance systems alert clinicians to the early signs of sepsis, enabling initiation of timely treatment. The

ideal surveillance system should be available in real-time at the point-of-care with the need for early identification of patients with sepsis strengthened by the known linear increase in the risk of mortality with each hour of delay in appropriate antibiotic treatment.^{18 19} The establishment of electronic health records systems to capture vital signs and laboratory values of patients have been shown to be very effective in alerting clinicians to possible clinical deterioration due to sepsis.²⁰ By harnessing these existing technologies, local and regional information system networks for sepsis could be established in Africa.

3. Invest in health workers

Existing data highlight a marked disparity in annual health expenditure per capita between LMICs and HICs; however, the economic cost of maternal and neonatal sepsis in Africa is not clearly defined.^{5 21} The WHA sepsis resolution advocates for well-coordinated efforts by politicians, policymakers, healthcare administrators, clinicians and researchers to reduce the impacts of sepsis.¹⁰ Understaffing and a higher workload is a common problem in health facilities across SSA⁹ and is linked to higher infection rates as well as adverse outcomes.^{22 23} Addressing this challenge will require accelerated progress in health worker recruitment and training.

Training and education of health workers, especially nurses and other frontline clinicians, is vital in improving the outcomes of critically ill patients.^{24 25} Retention of skilled staff should be a key consideration as 'brain drain' disproportionately affects health workers in SSA. Strategies should include improved remuneration, better conditions of service, task shifting and increased collaboration between those working in community and hospital settings. In SSA, mobile devices have been successfully used to train health workers.^{26 27} Aside from promoting contact with health workers in remote and hard-to-reach areas, electronic training has the added advantage of supporting provider-to-provider communication, feedback and professional networking. Programmes targeting education of health workers in prevention and early recognition and treatment of maternal and newborn sepsis should be prioritised.

4. Improve the early identification of sepsis in facilities

Several sepsis diagnostic scoring systems have been developed to identify changes in physiological parameters which precede patient deterioration²⁸ and alert clinicians to promptly initiate treatment. Examples of these tools for use in the general adult population include the Modified Early Warning Score, Universal Vital Assessment Score,²⁹ National Early Warning Score 2 and the quick Sepsis Organ Failure Assessment (qSOFA) score. Importantly, none of these scores have been validated in pregnant women or newborns. As such, they are unlikely to account for physiological differences in these population groups.³⁰

The Sepsis in Obstetrics Score, developed to detect maternal sepsis, appeared promising in initial reports from the USA; however, it was not superior to general sepsis scoring systems when tested in other regions.³¹

The obstetric modified SOFA score is an interesting adaptation of qSOFA that requires only clinical data for assessment and thus can be performed quickly and independently of laboratory tests.³² The UK Modified Early Obstetrics Warning Score (MEOWS) was developed for the obstetric population to aid the identification of clinical signs of deterioration in women developing critical illness. The MEOWS is easy to administer and has been shown to have a high sensitivity (89%) and reasonable specificity (79%) in predicting maternal morbidity, though it is not specific to sepsis.³³

The shock index (SI), defined as ratio of heart rate to systolic blood pressure, has recently been identified as a consistent predictor of adverse outcomes.^{34 35} Various thresholds have been proposed for use of SI as an early warning system. In a prospective study involving women with postpartum haemorrhage or sepsis, the SI was found to be a consistent predictor of adverse outcomes compared with conventional vital signs.³⁶

The introduction of maternal and neonatal sepsis scoring systems may be the first step towards achieving a reliable surveillance system for sepsis in SSA. Electronic surveillance tools which monitor physiological parameters should be developed to facilitate automatic recognition of the early signs of sepsis in health facilities.

5. Ensure laboratory diagnostics for sepsis are affordable and available

Infection is usually indicated by a rise in peripheral inflammatory makers such as the white blood cell count and C reactive protein. Cultures of relevant samples such as blood, urine, sputum and tissues provide microbiological confirmation of infections that give rise to sepsis. Rapid identification of pathogens from blood cultures has been shown to positively influence patient survival, length of hospitalisation and rational antibiotic prescriptions.³⁷ The development of local antibiograms is desirable to help facilities understand their local bacterial pathogens and match them to antibiotic choices. The establishment and maintenance of reference laboratories to support more sophisticated testing is essential.

In many LMIC settings, basic diagnostic modalities are lacking. Building local clinical microbiology laboratory capacity is an urgent priority and should incorporate regular training for staff to ensure they stay abreast with modern diagnostic methods. Recent research has focused on identifying biomarkers and molecular detection tools to increase the speed and reliability of diagnosis.³⁸ However, to ensure accessibility and affordability of modern diagnostic and biomarker testing, global health researchers and funders must ensure innovation in diagnostics and in their accessibility and affordability in LMIC settings.

Direct pathogen detection platforms promote the simultaneous identification of multiple bacterial and fungal species.^{38 39} While there are several rapid molecular diagnostic platforms to identify bacterial and viral pathogens commercially available, none are widely available or affordable for resource-poor settings. Although

rapid, innovative and low-cost testing devices will improve the identification of causal micro-organisms; to improve outcomes, this must be matched with availability of appropriate antimicrobial agents. This requires certainty of supply chains, committed funding, along with strategies at institutional level for antimicrobial stewardship to prevent the emergence of resistant organisms.

6. Improve quality of care through sepsis specific guidelines

While adherence to protocols and clinical guidelines can improve survival and outcomes in sepsis, in a survey of 185 hospitals across SSA, only 1.5% used clinical sepsis guidelines. Lack of basic facilities, equipment and medicines were identified as barriers to implementation.⁷ Resource limitations profoundly impact the ability to implement internationally recommended sepsis guidelines in SSA.⁴⁰

In high-income settings, significant reductions in maternal and neonatal sepsis have been achieved through quality improvement initiatives to promote adherence to evidenced-based management of sepsis.^{41 42} Quality improvement initiatives include educational programmes to increase awareness of the signs of sepsis; use of triage-based algorithms and decision support tools to screen for sepsis,^{43 44} use of treatment order-sets⁴⁵ and initiatives to reduce time to antibiotics.⁴⁶

Prompt administration of antibiotics remains the cornerstone of sepsis management. In the face of a growing burden of antimicrobial resistance and antibiotic-associated infections such as *Clostridioides difficile* infection, a greater alignment between antimicrobial stewardship and sepsis prevention and treatment programmes is essential.⁴⁷ Complementary strategies such as de-escalation protocols and prompt discontinuation of antibiotics when no longer indicated should be incorporated into sepsis quality improvement initiatives. Little is known about antimicrobial resistance in maternity units globally⁴⁸ and this information is essential to guide policy on the safe administration of prophylactic antibiotics during childbirth.

Another obvious gap is found in the Third International Consensus on Sepsis (Sepsis-3)¹ which does not provide criteria for diagnosis of sepsis in pregnant women. The ability of clinicians in SSA countries to adhere to sepsis guidelines is also limited by shortage of appropriately trained health workers, equipment and other supporting infrastructure.⁷

7. Increase access and use of vaccines to prevent the most common infections that lead to sepsis in pregnancy and the newborn period

Vaccinations are estimated to prevent 2–3 million infection-associated deaths every year among women and children.⁴⁹ Vaccines also have an indirect action on minimising antimicrobial resistance by reducing febrile illnesses and antibiotic use.⁵⁰ A vaccine for Group B Streptococcus is currently under evaluation and if effective, will have the potential to save 150 000 newborn lives every year and prevent up to 260 000 cases of maternal and

newborn sepsis.⁵¹ Seasonal influenza caused by types A, B, C and D of influenza viruses is a public health problem which can be prevented by vaccination. Although the burden of seasonal influenza in Africa is yet to be fully described, the morbidity, mortality and economic losses from this infection are likely to be very high.⁵²

Despite widespread availability of whole-cell pertussis (wP) vaccines, the burden of pertussis remains unacceptably high in the African continent.⁵³ This has been attributed to factors such as a lack of disease awareness and surveillance, diagnostic limitations and low prioritisation of this disease by policy makers. Intensified efforts to promote greater coverage with three primary doses of diphtheria-tetanus-pertussis vaccines and booster doses in older children and adolescents in Africa is also required.

Widespread protocols around immunisation for both adults and children is likely to decrease sepsis mortality in SSA. Current barriers to vaccination in this region include armed conflict, funding shortfalls, vaccine shortages and a lack of political will.⁵⁴ Measures to address these barriers include the establishment of lasting peace across the continent, improved funding for vaccination programmes and strengthening regional supply chains.

8. Strengthen infection prevention and control programmes and best practices during labour, childbirth and postnatal care

Water, sanitation and hygiene (WASH) is critical for the prevention of sepsis and provision of safe and quality care.⁵⁵ Less than 30% of delivery rooms in four East African countries reported access to a safe water supply⁵⁶ while a global review of health facilities in low-resource settings found that 40% had no water supply and 19% lacked adequate sanitation conditions.⁶ Improvements in WASH could reduce the total burden of disease worldwide by as much as 10%.⁵⁷ Implementation of basic measures such as hand hygiene, reducing overcrowding in maternity and neonatal units, promoting hygienic care of the umbilical cord and kangaroo care for small and/or preterm babies have all been shown to mitigate the risk of infection.^{58 59} Thus, the provision of safe water and promotion of effective hygiene practices should be emphasised and urgently prioritised in healthcare settings in SSA. This could be achieved by securing the commitment of policy makers to provide clean water in health facilities and by educating health workers and the wider population on the importance of adhering to standard hygiene practices.

9. Undertake high-quality research in pregnant and newborn African populations

There are imminent dangers in transferring effective solutions for sepsis management in HICs to LMICs. This is mirrored by recent well designed randomised controlled trials which have reflected worse outcomes for both children and adults with sepsis treated with

fluid boluses in Africa.^{13 60 61} The two trials conducted in Zambia comparing an early goal directed therapy (EGDT) protocol (involving intravenous fluids among other measures) to usual care found increased in-hospital mortality in the EGDT arm compared with usual care.^{13 61} High-quality research led by local investigators needs to be conducted to provide answers to conundrums associated with sepsis management among pregnant and newborn populations. These include determining the ideal type and volume of resuscitation fluid for sepsis, ascertaining the role of targeted/precision medicine approaches, rapid diagnostic tests and defining predictors of sepsis long-term morbidity and mortality.

CONCLUSION

Maternal and neonatal sepsis is under-recognised in SSA with significant mortality and morbidity consequences. Although mostly preventable, a complex interplay of several factors is hampering a robust response to the threat posed by sepsis in SSA.

A multimodal approach is required to tackle maternal and neonatal sepsis within the fragile health systems of the SSA region. Pragmatic steps in alignment with the WHA sepsis resolution include increasing public and health worker awareness of sepsis, establishing surveillance networks, developing laboratory and human capacity, facilitating research and the development of clinical guidelines in addition to promoting routine vaccination and good infection prevention and control practices.

Contributors SY and AO conceptualised and wrote the initial draft. EFS, JH, KT and KW provided comments and edits. All authors commented on the first draft and signed off on the final version.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement There are no data in this work.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID ID

Sanni Yaya <http://orcid.org/0000-0002-4876-6043>

REFERENCES

- Singer M, Deutschman CS, Seymour CW, *et al*. The third International consensus definitions for sepsis and septic shock (Sepsis-3). *JAMA* 2016;315:801–10.
- Rudd KE, Johnson SC, Agesa KM, *et al*. Global, regional, and national sepsis incidence and mortality, 1990–2017: analysis for the global burden of disease study. *Lancet* 2020;395:200–11.
- Say L, Chou D, Gemmill A, *et al*. Global causes of maternal death: a who systematic analysis. *Lancet Glob Health* 2014;2:e323–33.
- Lawn J, Mongi P, Cousens S. Opportunities for Africa's Newborns Practical data, policy and programmatic support for newborn care in Africa, 2006. Available: <https://www.who.int/pmnch/media/publications/oanfullreport.pdf>
- Yaya S, Bishwajit G, Okonofua F, *et al*. Under five mortality patterns and associated maternal risk factors in sub-Saharan Africa: a multi-country analysis. *PLoS One* 2018;13:e0205977.
- World Health Organization 2015. *Water, sanitation and hygiene in health care facilities: status in low- and middle-income countries and way forward*. Geneva: World Health Organization, 2015.
- Baelani I, Jochberger S, Laimer T, *et al*. Availability of critical care resources to treat patients with severe sepsis or septic shock in Africa: a self-reported, continent-wide survey of anaesthesia providers. *Crit Care* 2011;15:R10.
- World Health Organization. Maternal sepsis, 2018. Available: <https://www.who.int/reproductivehealth/maternal-sepsis/en/>
- Griffiths P, Renz A, Hughes J, *et al*. Impact of organisation and management factors on infection control in hospitals: a scoping review. *J Hosp Infect* 2009;73:1–14.
- World Health Organization 2017. *Seventieth World Health Assembly. WHA Resolution A70/13 - Improving the prevention, diagnosis and clinical management of sepsis. Report by the Secretariat*. Geneva: World Health Organization, 2017.
- Rubulotta FM, Ramsay G, Parker MM, *et al*. An international survey: public awareness and perception of sepsis. *Crit Care Med* 2009;37:167–70.
- Lewis JM, Feasey NA, Rylance J. Aetiology and outcomes of sepsis in adults in sub-Saharan Africa: a systematic review and meta-analysis. *Crit Care* 2019;23:212.
- Andrews B, Semler MW, Muchemwa L, *et al*. Effect of an early resuscitation protocol on in-hospital mortality among adults with sepsis and hypotension: a randomized clinical trial. *JAMA* 2017;318:1233–40.
- Vincent J-L. Increasing awareness of sepsis: world sepsis day. *Crit Care* 2012;16:152.
- Broadband Commission for Sustainable Development. Digital health: a call for government leadership and cooperation between ICT and health. broadband Commission for sustainable development, 2017. Available: <http://vitalwave.com/wp-content/uploads/2017/02/Broadband-Commission-Digital-Health-Final-Report-20170222.pdf> [Accessed 01 Dec 2019].
- Jabaley CS, Blum JM, Groff RF, *et al*. Global trends in the awareness of sepsis: insights from search engine data between 2012 and 2017. *Crit Care* 2018;22:7.
- Agarwal S, Perry HB, Long L-A, *et al*. Evidence on feasibility and effective use of mHealth strategies by frontline health workers in developing countries: systematic review. *Trop Med Int Health* 2015;20:1003–14.
- Kumar A, Roberts D, Wood KE, *et al*. Duration of hypotension before initiation of effective antimicrobial therapy is the critical determinant of survival in human septic shock. *Crit Care Med* 2006;34:1589–96.
- Han YY, Carcillo JA, Dragotta MA, *et al*. Early reversal of pediatric-neonatal septic shock by community physicians is associated with improved outcome. *Pediatrics* 2003;112:793–9.
- Makam AN, Nguyen OK, Auerbach AD. Diagnostic accuracy and effectiveness of automated electronic sepsis alert systems: a systematic review. *J Hosp Med* 2015;10:396–402.
- Rudd KE, Kissoon N, Limmathurotsakul D, *et al*. The global burden of sepsis: barriers and potential solutions. *Crit Care* 2018;22:232.
- Rogowski JA, Staiger D, Patrick T, *et al*. Nurse staffing and NICU infection rates. *JAMA Pediatr* 2013;167:444–50.
- Küng E, Waldhör T, Rittenschöber-Böhm J, *et al*. Increased nurse workload is associated with bloodstream infections in very low birth weight infants. *Sci Rep* 2019;9:6331.
- He Z, Bishwajit G, Zou D, *et al*. Burden of common childhood diseases in relation to improved water, sanitation, and hygiene (wash) among Nigerian children. *Int J Environ Res Public Health* 2018;15:pii: E1241.
- Nguyen HM, Schiavoni A, Scott KD, *et al*. Implementation of sepsis management guideline in a community-based teaching hospital - can education be potentially beneficial for septic patients? *Int J Clin Pract* 2012;66:705–10.
- O'Donovan J, Bersin A. Controlling Ebola through mHealth strategies. *Lancet Glob Health* 2015;3:e22.
- Otu A, Ebenso B, Okuzu O, *et al*. Using a mHealth tutorial application to change knowledge and attitude of frontline health workers to Ebola virus disease in Nigeria: a before-and-after study. *Hum Resour Health* 2016;14:5.
- Goldhill DR, McNarry AF. Physiological abnormalities in early warning scores are related to mortality in adult inpatients. *Br J Anaesth* 2004;92:882–4.
- Johnson J, Akinboyo IC, Curless MS, *et al*. Saving neonatal lives by improving infection prevention in low-resource units: tools are needed. *J Glob Health* 2019;9:010319.

- 30 Albright CM, Has P, Rouse DJ, *et al*. Internal validation of the sepsis in obstetrics score to identify risk of morbidity from sepsis in pregnancy. *Obstet Gynaecol* 2017;130:747–55.
- 31 Aarvold ABR, Ryan HM, Magee LA, *et al*. Multiple organ dysfunction score is superior to the Obstetric-Specific sepsis in obstetrics score in predicting mortality in septic obstetric patients. *Crit Care Med* 2017;45:e49–57.
- 32 Bowyer L, Robinson HL, Barrett H, *et al*. SOMANZ guidelines for the investigation and management sepsis in pregnancy. *Aust N Z J Obstet Gynaecol* 2017;57:540–51.
- 33 Cook CA. Implementing the modified early obstetric warning score (MEOWS) to detect early signs of clinical deterioration and decrease maternal mortality. *J Obst Gyn Neo* 2014;43:S22
- 34 Nathan HL, El Ayadi A, Hezelgrave NL, *et al*. Shock index: an effective predictor of outcome in postpartum haemorrhage? *BJOG* 2015;122:268–75. -.
- 35 El Ayadi AM, Nathan HL, Seed PT, *et al*. Vital sign prediction of adverse maternal outcomes in women with hypovolemic shock: the role of shock index. *PLoS One* 2016;11:e0148729.
- 36 Nathan HL, Seed PT, Hezelgrave NL, *et al*. Shock index thresholds to predict adverse outcomes in maternal hemorrhage and sepsis: a prospective cohort study. *Acta Obstet Gynaecol Scand* 2019;98:1178–86.
- 37 Beekmann SE, Diekema DJ, Chapin KC, *et al*. Effects of rapid detection of bloodstream infections on length of hospitalization and hospital charges. *J Clin Microbiol* 2003;41:3119–25.
- 38 Al Jalbout N, Troncoso R, Evans JD, *et al*. Biomarkers and molecular diagnostics for early detection and targeted management of sepsis and septic shock in the emergency department. *J Appl Lab Med* 2019;3:724–9.
- 39 De Angelis G, Posteraro B, De Carolis E, *et al*. T2Bacteria magnetic resonance assay for the rapid detection of ESKAPEc pathogens directly in whole blood. *J Antimicrob Chemother* 2018;73:iv20–6.
- 40 Ndadane N, Maharaj RC. The epidemiology of sepsis in a district hospital emergency centre in Durban, KwaZulu natal. *Afr J Emerg Med* 2019;9:123–6.
- 41 Paul R, Melendez E, Stack A, *et al*. Improving adherence to PaIS septic shock guidelines. *Pediatrics* 2014;133:e1358–66.
- 42 Paul R, Neuman MI, Monuteaux MC, *et al*. Adherence to PaIS sepsis guidelines and hospital length of stay. *Pediatrics* 2012;130:e273–80.
- 43 Chamberlain DJ, Willis E, Clark R, *et al*. Identification of the severe sepsis patient at triage: a prospective analysis of the Australasian triage scale. *Emerg Med J* 2015;32:690–7.
- 44 Gatewood Medley O'Keefe, Wemple M, Greco S, *et al*. A quality improvement project to improve early sepsis care in the emergency department. *BMJ Qual Saf* 2015;24:787–95.
- 45 McDonald CM, West S, Dushenski D, *et al*. Sepsis now a priority: a quality improvement initiative for early sepsis recognition and care. *Int J Qual Health Care* 2018;30:802–9.
- 46 Emerson BL, Ciaburri R, Brophy C, *et al*. Decreasing time to antibiotics for patients with sepsis in the emergency department. *Pediatr Qual Saf* 2019;4:e173.
- 47 Fitzpatrick F, Tarrant C, Hamilton V, *et al*. Sepsis and antimicrobial stewardship: two sides of the same coin. *BMJ Qual Saf* 2019;28:758–61.
- 48 Thompson K, Hirst J, Walker K. Maternal and newborn sepsis worldwide. *Sepsis Redefined* 2019;21.
- 49 World Health Organization. Fact sheet: immunization coverage, 2019. Available: <https://www.who.int/news-room/fact-sheets/detail/immunization-coverage> [Accessed 19 Nov 2019].
- 50 Klugman KP, Black S. Impact of existing vaccines in reducing antibiotic resistance: primary and secondary effects. *Proc Natl Acad Sci U S A* 2018;115:12896–901.
- 51 Seale AC, Bianchi-Jassir F, Russell NJ, *et al*. Estimates of the burden of group B streptococcal disease worldwide for pregnant women, stillbirths, and children. *Clin Infect Dis* 2017;65:S200–19.
- 52 Katz MA, Schoub BD, Heraud JM, *et al*. Influenza in Africa: uncovering the epidemiology of a long-overlooked disease. *J Infect Dis* 2012;206 Suppl 1:S1–4.
- 53 Muloiwira R, Wolter N, Mupere E, *et al*. Pertussis in Africa: findings and recommendations of the global pertussis initiative (GPI). *Vaccine* 2018;36:2385–93.
- 54 Nshimirimana D, Mihigo R, Clements CJ. Routine immunization services in Africa: back to basics. *Journal of Vaccines & Immunization* 2013;1:6–12.
- 55 Mullany LC, Darmstadt GL, Katz J, *et al*. Risk factors for umbilical cord infection among newborns of southern Nepal. *Am J Epidemiol* 2007;165:203–11.
- 56 Gon G, Restrepo-Méndez MC, Campbell OMR, *et al*. Who delivers without water? A multi country analysis of water and sanitation in the childbirth environment. *PLoS One* 2016;11:e0160572.
- 57 Prüss-Ustün A, Bartram J, Clasen T, *et al*. Burden of disease from inadequate water, sanitation and hygiene in low- and middle-income settings: a retrospective analysis of data from 145 countries. *Trop Med Int Health* 2014;19:894–905.
- 58 World Health Organization. Survive and thrive: transforming care for every small and sick newborn, 2019. Available: www.who.int/maternal_child_adolescent/documents/care-small-sick-newborns-survive-thrive/en/
- 59 World Health Organization. Who recommendations for prevention and treatment of maternal peripartum infections, 2015. Available: www.who.int/reproductivehealth/publications/maternal_perinatal_health/peripartum-infections-guidelines/en/
- 60 Maitland K, Kiguli S, Opoka RO, *et al*. Mortality after fluid bolus in African children with severe infection. *N Engl J Med* 2011;364:2483–95.
- 61 Andrews B, Muchemwa L, Kelly P, *et al*. Simplified severe sepsis protocol: a randomized controlled trial of modified early goal-directed therapy in Zambia. *Crit Care Med* 2014;42:2315–24.